

# 2021 Suisun Marsh Vegetation Map Update Solano County, California

**June  
2024**

**A Report to the California  
Department of Water Resources**

**Prepared by:**  
**Geographical Information Center**  
**North State Planning and Development Collective**  
**California State University, Chico**

**Malak Saleh, Biologist Interpreter**  
**Kristin Quigley, Lead Biologist Interpreter**  
**Jaclyn Deloach, Biologist Interpreter**  
**Erik Fintel, Project Manager**  
**Laura Askim, Biologist Interpreter**





## Executive Summary

This report summarizes the methods and results of the 2021 Suisun Marsh triennial vegetation map update. This update by the Geographical Information Center (GIC) of the North State Planning and Development Collective at California State University, Chico is part of an ongoing triennial vegetation monitoring program for the Suisun Marsh. The project tracks changes in the Suisun Marsh vegetation over time to fulfill specific permit requirements of the Suisun Marsh Plan of Protection of 1984, the Suisun Marsh Preservation Agreement of 1986, and the 2015 Suisun Marsh Preservation Agreement. This is the seventh update using the current mapping standards originally implemented in 1999. All of the previous vegetation maps from 1999 to 2018 can be viewed and downloaded using the online California Department of Fish and Wildlife (CDFW) Biogeographic Information and Observation System (BIOS); the links to the associated reports are included in the map metadata. For detailed information regarding the history and evolution of this project, see Appendix A in the 2012 update report ([Boul and Keeler-Wolf 2016](#)).

The final map covers 91,486 acres, containing vegetation and land use polygons ranging from 0.001 acres to 304 acres and averaging 1.7 acres. The tidally influenced acreage is covered by 6,665 vegetation polygons; 32,956 vegetation polygons cover the leveed acreage.

The primary and secondary Salt Marsh Harvest Mouse (SMHM) habitat types listed in the Bay Delta Conservation Plan (DWR 2013) were used to calculate the potential habitat for SMHM. The 2021 triennial vegetation update suggests there are 46,661 acres of potential habitat in Suisun Marsh, with approximately 81% of this habitat in leveed areas, and the remaining habitat in tidally influenced areas. The leveed areas have seen a 6.5% decrease in potential habitat since 2018, and an overall 9.2% decrease since 1999. The tidal areas of the Marsh have seen a 2.2% increase in potential SMHM habitat since 2018, with an overall 59.3% increase since 1999. Over the entire marsh, potential habitat has decreased by 5% since 2018 and 1% since 1999.

The distribution and spread of non-native plant species of concern were also analyzed across the entire marsh, as well as within each of the four management regions. The non-native species of concern include: *Arundo donax*, *Carpobrotus edulis*, *Cortaderia selloana*, *Eucalyptus* spp., *Lepidium latifolium*, *Phragmites australis*, and *Salsola soda*. Comparisons were made between current distributions and the original 1999 map and the previous 2018 map. The most substantial increase occurred in *Phragmites australis*, the most widespread non-native species of concern in the Marsh. *P. australis* has increased from 693 acres in 1999 to 4,296 acres in 2018, and 4,400 acres in 2021. Full change analysis and discussion of each species can be found further on in this report.

# Table of Contents

<b>INTRODUCTION .....</b>	<b>1</b>
<b>METHODS .....</b>	<b>4</b>
FIELD DATA COLLECTION .....	4
2021 AERIAL PHOTOGRAPH INTERPRETATION, HEADS-UP DIGITIZING, AND ATTRIBUTING .....	4
MAPPING RULES.....	4
MAP ATTRIBUTES .....	4
METHODS FOR ANALYSIS.....	5
ACCURACY ASSESSMENT .....	7
<b>RESULTS .....</b>	<b>8</b>
MAPPING .....	8
VEGETATION CHANGE ANALYSIS .....	8
<i>Salt Marsh Harvest Mouse Habitat</i> .....	9
<i>Non-Native Species of Concern</i> .....	12
<b>DISCUSSION, CONCLUSIONS, AND RECOMMENDATIONS.....</b>	<b>21</b>
<i>Salt Marsh Harvest Mouse Habitat</i> .....	21
<i>Change in Tidal Habitat</i> .....	23
<i>Non-Native Species of Concern</i> .....	24
<i>Changes in Slough Cover</i> .....	24
FUTURE MAPPING EFFORTS .....	24
<b>LITERATURE CITED .....</b>	<b>26</b>

## Tables

<b>Table 1: Total acreage and polygons mapped in the 2021 Suisun Map Update .....</b>	<b>8</b>
---	----------

## Figures

<b>Figure 1: Location of the Suisun Marsh study area. ....</b>	<b>1</b>
<b>Figure 2: A map showing the four geographic management regions (1-4) in the Suisun Marsh.....</b>	<b>7</b>
<b>Figure 3: The acreage and distribution for the potential Salt Marsh Harvest Mouse habitat .....</b>	<b>9</b>
<b>Figure 4: 2021 Salt Marsh Harvest Mouse habitat by region and habitat type (leveed or tidal) .....</b>	<b>10</b>
<b>Figure 5: Acreages of the less abundant non-native species of concern .....</b>	<b>12</b>
<b>Figure 6: Acreages of the more abundant non-native species of concern.....</b>	<b>13</b>
<b>Figure 7: <i>Arundo donax</i> distribution by management region .....</b>	<b>14</b>
<b>Figure 8: <i>Carpobrotus edulis</i> distribution by management region.....</b>	<b>15</b>
<b>Figure 9: <i>Cortaderia selloana</i> distribution by management region .....</b>	<b>15</b>
<b>Figure 10: <i>Eucalyptus</i> spp. distribution by management region.....</b>	<b>16</b>
<b>Figure 11: <i>Lepidium latifolium</i> distribution by management region.....</b>	<b>16</b>
<b>Figure 12: <i>Phragmites australis</i> distribution by management region.....</b>	<b>17</b>
<b>Figure 13: <i>Salsola soda</i> distribution by management region .....</b>	<b>17</b>
<b>Figure 14: Example of Bare Ground.....</b>	<b>22</b>
<b>Figure 15: Example of Levee Break.....</b>	<b>23</b>

## **Appendices**

**Appendix A:** Suisun Marsh Vegetation Classification and Hierarchy

**Appendix B:** Vegetation Type Descriptions in Suisun Marsh

**Appendix C:** Map Classes Used to Map Suisun Marsh

**Appendix D:** 2021 Vegetation Mapping Attributes and Descriptions

**Appendix E:** Salt Marsh Harvest Mouse Habitat by Vegetation Mapping Type

**Appendix F:** Acreage and Acreage Change of Salt Marsh Harvest Mouse Habitat

**Appendix G:** Acreage and Acreage Change of the Non-Native Species of Concern

**Appendix H:** Reconnaissance Field Form and Protocol

## Introduction

*Note: This introduction is credited to CDFW's Vegetation Classification and Mapping Program ([VegCAMP 2018](#)).*

The Suisun Marsh is located in Solano County, CA, and is part of the San Francisco Bay / Sacramento–San Joaquin River Delta estuary ecosystem (Figure 1). It is one of the largest contiguous brackish marshes remaining in the United States, covering over 69,000 acres of tidal and managed seasonal wetlands. This Marsh is a key wintering area for waterfowl and supports several sensitive plant and animal species. Since 1999, the Marsh has experienced significant variations in salinity conditions, including prolonged droughts. Plant communities can be expected to change over time due to these natural environmental factors, but also from direct management activities such as flooding regime changes, weed control, and plantings, as well as from accidental introductions of invasive weeds and levee breaches.



**Figure 1: Location of the Suisun Marsh study area.**

As part of the monitoring program in the Plan of Protection for Suisun Marsh and as required in the 1981 US Fish and Wildlife Service Biological Opinion, a Triennial Vegetation Survey is necessary to monitor changes in Salt Marsh Harvest Mouse (SMHM) habitat. The current methodology for this survey was adopted in 1999 ([Keeler-Wolf and Vaghti 2000](#)) and uses aerial photography in combination with ground verification to document the overall vegetation composition of the Marsh and to monitor SMHM habitat. The methodology documents the changes in preferred habitat for the SMHM and gathers vegetation information for a variety of other purposes.

This methodology followed the National Vegetation Classification (NVC) standards and resulted in a vegetation classification based on quantitative vegetation sampling (Appendix A). The vegetation classification was used to create vegetation type descriptions (Appendix B) and a mapping classification (Appendix C). While the mapping classification is based on the vegetation classification, the mapping classification is limited by what is discernable from the aerial imagery. It also includes mapping units that are not currently accepted NVC vegetation types and mapping units that represent land use or non-vegetated types. Since the 1999 map, there have been updates in 2000, 2003, 2006, 2009, 2012, 2015, and 2018. This report documents changes based on the 2021 vegetation map update.

As requested by the Department of Water Resources (DWR) and with CDFW Bay Delta Region agreement, for each map update, vegetation change has been analyzed for tidal wetlands, leveed wetlands, and marsh-wide. Tidal wetlands (including muted tidal wetlands) are those areas naturally affected by regular tidal fluctuation. These areas may or may not be vegetated with vascular or non-vascular plants, and may or may not have any evidence of human modification such as ditches, excavations, levees, or berms. The leveed-managed wetlands are those areas that are completely enclosed by exterior levees, with water application and drainage controlled through water management infrastructure.

The 2000 remap effort ([Vaghti and Keeler-Wolf 2001](#)) was an exploratory change detection study designed to define significant changes in vegetation in the Suisun Marsh ecosystem. Less than 1% of the polygons changed between June 16, 1999, and July 2, 2000. These minor changes included a net loss of 65 acres for *Salicornia pacifica* vegetation types, an 18-acre increase in vegetation dominated by *Lepidium latifolium*, and a 143-acre decrease in Annual Grasses. From this exploratory change detection, it was determined that the map update process would occur every three years.

The 2003 remap effort ([Vaghti and Keeler-Wolf 2004](#)) showed a 16.8% change in the vegetation across the entire study area since the 1999 product. Medium Wetland Graminoids, *Bolboschoenus maritimus*, Short Wetland Herbs, Medium Wetland Herbs, and *Bolboschoenus maritimus* – *Salicornia pacifica* were the five types with the greatest increase in acreage. *Distichlis spicata*, *Salicornia*, *Distichlis spicata* – Annual Grasses, *Distichlis spicata* – *Salicornia pacifica*, and Open Water were the five types with the greatest decrease in acreage over the study period. Also determined was a 16.7% change in leveed wetland vegetation and a 17.2% change in tidal wetland vegetation.

The 2006 remap study ([Boul and Keeler-Wolf 2008](#)) used the 1999 vegetation map as the baseline and followed the 2000 and 2003 change detection methodology. Several vegetation changes found in the 2006 update were of note: 1) a 174% increase in flooded wetlands (due to severe storms resulting in levee breaches), 2) a net loss of 945 acres of *Salicornia pacifica* vegetation types since 1999, 3) a net gain of 780 acres of *Phragmites australis* since 1999, 580 acres of which established since 2003, and 4) acreage decrease or stabilization of several of the non-native species of concern. Several issues with the remapping process and change detection protocol emerged and VegCAMP suggested changes to the protocol. These were implemented in the 2009 vegetation remap.

The 2009 remap study ([VegCAMP 2012](#)) used an updated protocol developed to accommodate advances in available technology and mitigate past inconsistencies; polygon modification was discontinued, and



polygons were instead mapped anew, without using previous linework. The protocol update is discussed in detail in Appendix A in [Boul and Keeler-Wolf \(2016\)](#). The 2009 remap showed that potential Salt Marsh Harvest Mouse habitat (*Salicornia pacifica* dominated vegetation) had increased since 1999 and that two non-native species of concern, *Phragmites australis*, and *Lepidium latifolium*, were still increasing within the Marsh. Interestingly, in the leveed areas of the Marsh, both *Phragmites australis* and *Salicornia pacifica* vegetation seemed to be increasing the most where there had been open water in 2006.

The 2012 remap study ([Boul and Keeler-Wolf 2016](#)) showed a marsh-wide 7% increase in potential habitat for SMHM since 2009 and a 1% increase since 1999. By 2012, *Phragmites australis* had increased threefold in acreage since the first map in 1999. The increase was and has been presumed to be in the non-native genotype. Stands dominated by *Lepidium latifolium* showed a marsh-wide decrease in total acreage since 1999 but showed a 63% increase in the tidal habitats of the Marsh.

The 2015 remap study ([VegCAMP 2018](#)) showed a marsh-wide 3.9% decrease in potential habitat for SMHM since 2012 and a 7.3% decrease since 2009. From 2012 to 2015, the non-native *Phragmites australis* increased 18.5% marsh-wide, with a total increase of 325% (2,254 acres) since 1999. *Lepidium latifolium* acreage increased by 45% from 2012 to 2015. This remap also evaluated the potential use of satellite imagery instead of high-resolution imagery acquired by DWR for the mapping effort but determined that the lower-resolution satellite imagery taken at an uncontrolled time of day, year, and tidal stage made it unsuitable for the purposes of the triennial map update.

The 2018 remap study ([GIC 2022](#)) found a 12.4% increase in SMHM habitat since 2015 and a 4.2% increase since 1999. The study found a 46% increase in invasive *Phragmites australis* cover throughout the entire marsh from 2015 to 2018, and a total increase of 520% (3,603 acres) since 1999. *Lepidium latifolium* acreage increased by 90% from 2015 to 2018 but experienced an overall 68% decrease in cover since 1999.

This 2021 remap has shown an overall decrease in SMHM habitat since 2018. Overall, there has been a decrease of 2,444 acres (a 5% decrease) with a majority observed in leveed areas, where there was a 2,640-acre decrease, which reflects a 6.6% decrease in leveed habitat. The amount of tidal habitat increased by 196 acres and 2.23%. The findings of this remap indicate there has been a 0.9% decrease in habitat since 1999.



## Methods

### Field Data Collection

This mapping cycle included the collection of 309 reconnaissance points as well as 100 relevés that were collected from June through September of 2021. Map interpreters use these points to discern vegetative signatures and guide mapping type determinations of map polygons. The field data collection form and protocol can be found in Appendix H.

### 2021 Aerial Photograph Interpretation, Heads-up Digitizing, and Attributing

To create the 2021 Suisun Marsh vegetation map, vegetation was interpreted from a mosaic of the true color imagery that was flown in June 2021. Polygons were delineated using heads-up digitizing (i.e., a photo interpreter manually drew polygons around each stand of vegetation) in Esri's ArcGIS Pro 3.1.2, and polygon attributes were recorded within a file geodatabase.

All attributes were interpreted using the Suisun Marsh 2021 imagery as the base imagery. The photo interpreters obtained information primarily from the 2018 map and 2021 reconnaissance points, which were used during mapping to determine vegetative signatures and the appropriate mapping type for each polygon. Several other imagery sources were used as ancillary data, including 2021 NAIP, 2021 NAIP Color Infrared, all imagery available through Google Earth (including street view), and the 2018 NAIP imagery.

### Mapping Rules

Minimum mapping unit (MMU): Typically, the minimum mapping size is 0.25 acres. However, the photo interpreters use their best judgment to determine if a stand below 0.25 acre should be separately delineated. For example, a smaller polygon would be appropriate for any new visible occurrence of a non-native species of concern, such as *Phragmites australis*, *Arundo donax*, *Carpobrotus edulis*, *Eucalyptus* spp., and *Lepidium latifolium*.

Minimum mapping width: There are many long and narrow polygons within the Suisun Marsh study area, most of which are roads, ditches, levees, and sloughs. The minimum mapping width is typically 10 feet; however, if small sections of a stand fell below the minimum width, the polygon was not split.

### Map Attributes

The vegetation mapping classification used for this project is based on the vegetation classification created in 1999 (Keeler-Wolf and Vaghti 2000). The names of the mapping units have been updated over the years to reflect current nomenclature and/or current understanding of vegetation classification. We used the same map classes for the 2021 map that were used for the 2015 and 2018 maps with the addition of two new mapping units (*Spartina foliosa* Herbaceous Alliance and the California Vernal Pool and Grassland Matrix Group), which had not been used in previous mapping efforts. The *Spartina foliosa* Herbaceous Alliance was added as a result of 2021 reconnaissance survey

data, as this species had not been surveyed in past mapping efforts. The California Vernal Pool and Grassland Matrix Group was added as a result of mapper experience mapping this group in similar environments, and its identification was aided by 2005 mapping data provided by the California Department of Fish and Wildlife. Mappers visually verified the vernal pool signature to account for changes that have occurred since the vernal pool complex map was created. See Appendix C for a list of all the mapping types.

In addition to the mapping unit, several other attributes are assigned to each polygon. These attributes include percent vegetation cover, average vegetation height, degree of human disturbance, relative cover of invasive species, and habitat type (tidal versus leveed). For more information about each attribute, see Appendix D.

### Methods for Analysis

As previously mentioned, DWR and CDFW Bay Delta Region have requested that change analysis be done for tidal wetlands, leveed (managed) wetlands, and marsh-wide. To determine the areas within the marsh that are tidally influenced versus areas that are leveed (or managed), the 2018 remap habitat attribute was used as a reference. This was determined originally from a shapefile created by CDFW in 2008 ([VegCAMP 2018](#)). For the current remap, the polygons that were contained completely within areas that were called tidal habitat in 2018 were given the tidal habitat attribute again, as tidal areas tend to remain tidal. Polygons that were completely within the leveed habitat in 2018 were given the leveed habitat attribute after confirmation by a photo interpreter that the general area was still leveed. Additionally, the map was reviewed for levee breaks or levee additions that may have altered the habitat attribute. Sloughs were attributed separately and were not given a habitat type.

The vegetation map was checked for changes in stand-forming non-native species of concern between 1999 and 2021 and between 2018 and 2021. The analysis was performed marsh-wide (total area, leveed habitats, and tidal habitats) and within the four management regions (total area, leveed habitats, and tidal habitats). The non-native species of concern are as follows: *Arundo donax*, *Carpobrotus edulis*, *Cortaderia selloana*, *Eucalyptus* species, *Lepidium latifolium*, *Phragmites australis* (predominantly the non-native strain), and *Salsola soda*. These species are represented by the following mapping units:

- Arundo donax* Herbaceous Semi-Natural Association
- Carpobrotus edulis* or Other Ice Plants Herbaceous Semi-Natural Alliance
- Cortaderia (jubata, selloana)* Herbaceous Semi-Natural Alliance
- Eucalyptus (globulus, camaldulensis)* Woodland Semi-Natural Alliance
- Lepidium latifolium* Herbaceous Semi-Natural Alliance
- Phragmites australis* Herbaceous Alliance
- Salsola soda* Mapping Unit


At the time of the 2000 and 2003 Suisun Marsh vegetation change detection, less specific information was known about the habitat requirements for the protected Salt Marsh Harvest Mouse (*Reithrodontomys raviventris halicoetes*) in Suisun Marsh. Ten *Salicornia pacifica* (or “pickleweed”)

vegetation types or mapping units were collectively considered important habitat for the Salt Marsh Harvest Mouse (SMHM) in 2003 (Vaghti and Keeler-Wolf 2004). Since then, biologists have gained a better understanding of the habitat requirements of the SMHM and have considerably broadened the definition of what is considered SMHM habitat. The vegetation types that are considered SMHM habitat were updated in both 2013 and 2015. This report and the previous 2015 report use the 2013 habitat types. The updated 2015 habitat types include designating low-, medium-, and high-quality habitat, and will be beneficial to use in future reports, although the change in what is considered habitat will affect true change analysis for future remaps. See Appendix E for the full list of mapping units that are considered potential SMHM habitat in this report.

Vegetation change detection analysis for SMHM habitat and non-native species of concern was performed marsh-wide (separately for the total area, leveed areas, and tidal areas) and for each of the four management regions shown in Figure 2 (separately for the total area, leveed areas, and tidal areas). The percent change (acreage) was calculated using the following formula:

$$\frac{(\text{Current Year Acreage} - \text{Previous Acreage})}{\text{Previous Acreage}} * 100$$

See Appendices F and G for vegetation change in the SMHM habitat and non-native species of concern.



**Figure 2: A map showing the four geographic management regions (0-4) in the Suisun Marsh.**

'0' refers to areas outside of a defined management region.

## Accuracy Assessment

For this remap, an Accuracy Assessment was not performed due to delays in between the imagery flight date (June 2021) and the completion of the map (April 2024). Ideally, due to the fluctuating nature of the Marsh's vegetation, an Accuracy Assessment should be conducted no later than one year after the imagery was taken. A thorough Accuracy Assessment was conducted for the 2015 remap. For detailed methodology and results see [VegCAMP 2018](#).

## Results

### Mapping

The mapped project area in 2021 was 91,486 acres, covering the same area as the 2018 mapping effort. The Suisun Bay made up 24.7% of this area, tidal sloughs made up 3.1%, and areas mapped as vegetation or land use (such as roads or development) made up 72.2%. Of the vegetation and land use area, 16.3% was tidally influenced and the rest was leveed.

The current map consists of 39,643 polygons, while the 2018 map had 39,566 polygons. The total mapped acreage of vegetation changes slightly with each map update due to the tide level at the time the imagery was acquired and channel erosion or accretion since the area was last mapped.

**Table 1:** Total acreage and polygons mapped in the 2021 Suisun Map Update. This table is based on the values in the “habitat” attribute.


Area mapped as:	Acres	Polygons
Suisun Bay	22,627.45	1
Sloughs	2,797	22
Tidal vegetation/land use	11,206	6,664
Leveed vegetation/land use	54,854.86	32,956
<b>Total project area</b>	<b>91,486</b>	<b>39,643</b>

### Vegetation Change Analysis

The changes in Salt Marsh Harvest Mouse habitat and non-native species of concern were analyzed marsh-wide from 2018 to 2021. Changes from 1999 to 2021 are also discussed, and data from the 2015 map study were included for comparison purposes. These year-to-year comparisons are straightforward, as the total area designated as Suisun Marsh has remained static from 1999 to 2021. SMHM habitat and non-native species changes were also analyzed within each of the four management regions.

## Salt Marsh Harvest Mouse Habitat

### Marsh-wide



**Figure 3: The acreage and distribution across the four management regions for the potential Salt Marsh Harvest Mouse habitat mapped within the Suisun Marsh triennial vegetation maps in 1999, 2015, 2018, and 2021.**

Of the 66,062 acres of vegetation/land use (total area excluding sloughs and Suisun Bay) mapped in the Marsh in 2021, 71% (46,662 acres) is considered potential SMHM habitat, an overall decrease of 5% since 2018 (Figure 3). SMHM habitat comprises 80% of the tidal vegetation/land use areas and 69% of the leveed areas.


Excluding the non-native *Phragmites australis*, which has been used by SMHM and is considered low-quality habitat, there are 42,221 acres of habitat throughout the Marsh. In 2018, *Phragmites* accounted for 9% of the total SMHM habitat, and in 2021 it accounted for 9.5%.

There are 9,006 acres of SMHM habitat in tidal areas throughout the marsh, an increase of 2.2% since 2018 and 59% since 1999 (Appendix F). *Schoenoplectus (acutus, californicus)* mapping units (including *Schoenoplectus* mixed with *Rosa*, wetland herbs, and *Typha*) account for 3,804 acres of tidal habitat (42%, compared to 41% in 2018). *Phragmites* accounts for 1,910 acres of tidal habitat (remaining at 21% since 2018). *Distichlis* mapping units (including *Distichlis* mixed with annual grasses, *Cotula*, *Juncus*, *Salicornia*, and *Schoenoplectus americanus*) account for 1,286 acres of tidal habitat (14%, compared to 15% in 2018). *Schoenoplectus americanus* mapping units (including when mixed with *Lepidium*) account

for 1,377 acres of tidal habitat (remaining at 15% since 2018). These four vegetation types (*Schoenoplectus acutus/californicus*, *Phragmites*, *Distichlis*, and *Schoenoplectus americanus*) account for 93% of all tidal habitat acreage in 2021, an increase of 1% since 2018.

There are 37,656 acres of SMHM habitat in leveed areas throughout the marsh, a decrease of 6.6% since 2018 and a decrease of 9% since 1999 (Appendix F). *Salicornia pacifica* mapping units (including *Salicornia* mixed with annual grasses, *Cotula*, *Crypsis*, and *Sesuvium*) account for 14,994 acres of leveed habitat (40% compared to 39% in 2018). *Distichlis* mapping units account for 6,918 acres of leveed habitat (remaining at 18% since 2018). Mediterranean California Naturalized Annual and Perennial Grassland accounts for 5,845 acres of leveed habitat (16%, compared to 17% in 2018). The newly implemented California Vernal Pool and Grassland Matrix Group accounts for 788 acres of leveed habitat, or 2% of total leveed habitat. *Phragmites* accounts for 2,530 acres of leveed habitat (7%, compared to 6% in 2018). *Schoenoplectus (acutus, californicus)* mapping units account for 2,208 acres of leveed habitat (remaining at 6% since 2018). These six vegetation types (*Salicornia*, *Distichlis*, grassland, vernal pools, *Phragmites*, and *Schoenoplectus acutus/californicus*) account for 88% of all leveed habitat acreage in 2021 (86% in 2018).

#### By management region



**Figure 4: 2021 Salt Marsh Harvest Mouse habitat by region and habitat type (leveed or tidal)**



**Region 0:**

Management Region 0 covers 25,024 acres, the vast majority of which are open water (21,568 acres) and slough cover (2,443 acres) (Appendix F). The boundaries of this region contain just 3.73 acres of leveed SMHM habitat and 170 acres of tidal habitat (Figure 4). Tidal habitat in this region has increased by 115% since 2018 when there were just 79 acres of tidal habitat in this region. *Schoenoplectus (acutus, californicus)* mapping units occupy 140 acres (82%) of the tidal habitat in this region. *Phragmites* accounts for 14 acres (8%) of the habitat in this region.

**Region 1:**

Management Region 1 covers 18,988 acres. This region contains 12,071 acres (26%) of the SMHM habitat in the Marsh (Figure 3). Habitat in this region has decreased by 5% (647 acres) since 2018, when there were 12,718 acres. 78% of the SMHM habitat of this region is in the leveed areas (9,388 acres), where it has decreased by 712 acres (6%) since 2018 (Appendix F). In the tidal areas, habitat has seen an extremely slight decrease from 2,684 acres in 2018 to 2,682 acres in 2021 (a change of just two acres total, and no substantial percentage change). Total SMHM habitat has decreased by 9% since 1999 in this region. *Phragmites* accounts for 979 acres (8%) of the habitat in this region (Appendix G).

**Region 2:**

Management Region 2 covers 9,958 acres. This region contains 7,635 acres (16%) of the SMHM habitat in the Marsh (Appendix F). Habitat has decreased by 6% (477 acres) since 2018, when there were 8,112 acres. 72% of the SMHM habitat of this region is in the leveed areas (5,474 acres), where it has decreased by 505 acres (8%) since 2018 (Figure 4). In the tidal areas, habitat has increased from 2,137 acres in 2018 to 2,160 acres in 2021 (a 1% increase). Total SMHM habitat has increased by 10% since 1999 in this region (Appendix F). *Phragmites* accounts for 551 acres (7%) of the habitat in this region (Appendix G).

**Region 3:**

Management Region 3 covers 4,720 acres. This region contains 3,213 acres (7%) of the SMHM habitat in the Marsh (Figure 3). Habitat has decreased by 3% (93 acres) since 2018, when there were 3,307 acres (Appendix F). 76% of SMHM habitat of this region is in the leveed areas (2,456 acres), where it has decreased by 192 acres (7%) since 2018 (Figure 4). In the tidal areas, habitat has increased from 652 acres in 2018 to 757 acres in 2021 (a 14% increase). Total SMHM habitat has decreased by 10% since 1999 in this region (Appendix F). *Phragmites* accounts for 225 acres (7%) of the habitat in this region (Appendix G).

**Region 4:**

Management Region 4 covers 32,796 acres. This region contains 23,544 acres (50%) of the SMHM habitat in the marsh (Figure 3). Habitat has decreased by 5% (1,343 acres) since 2018, when there were 24,887 acres (Appendix F). 86% of the SMHM habitat of this region is in the leveed areas (20,307 acres), where it has decreased by 1,312 acres (6%) since 2018. In the tidal areas, habitat has decreased from 3,258 acres in 2018 to 3,237 acres in 2021 (a decrease of less than 1%). Total SMHM habitat has

increased by less than 1% since 1999 in this region. *Phragmites* accounts for 2,671 acres (11%) of the habitat in this region (Appendix G).

### Non-Native Species of Concern

All of the results of change detection for non-native species of concern are presented in Appendix G and are summarized or highlighted in the figure and discussion below.

#### Marsh-wide





Figure 5: Acreages of the less abundant non-native species of concern between 2015, 2018, and 2021



**Figure 6: Acreages of the more abundant non-native species of concern between 2015, 2018, and 2021**

*Arundo donax* has decreased by 1 acre since 2018 marsh-wide (Figure 5). There are currently 9 total acres; 5 acres are in leveed areas (a 24% decrease since 2018) and 4 acres are in tidal areas (a 14% increase since 2018). In 1999, there were 4.7 acres total throughout the marsh. The increase in *Arundo donax* cover since 1999 can be partially explained by improvements in imagery and mapping interpretation, as this species may not have been correctly identified in previous years. There is no minimum mapping unit (MMU) for invasive species, meaning there are mapped polygons of *Arundo donax* that cover less than .25 acres.

*Carpobrotus edulis* increased from 7.0 acres in 1999 to 40.2 acres in 2021 (Figure 5). The cover of *Carpobrotus edulis* has remained steady at 40.2 acres since 2018, and *Carpobrotus* does not appear to be spreading. However, this species has seen a 474% increase since mapping began in 1999 (Appendix G). Similarly to *Arundo donax*, the increase in acres is partially due to better photo interpretation and the lack of an MMU for invasive species in the later maps.

*Cortaderia selloana* has decreased from 10 acres in 2018 to 4.4 acres in 2021, falling below the 6 acres mapped in 2015 (Figure 5). 3.8 acres occur in leveed areas (a 34% decrease since 2018) and the remaining .6 acres are in tidal areas (an 84% decrease since 2018).

Mapped stands of *Eucalyptus* spp. have increased from 208.1 acres in 1999 to 320 acres in 2021 for an increase of 53.8%, with no observed increase from 2018 to 2021 (Appendix G). 306 of the 320 acres occur in leveed areas.


In 2018, there were 205.7 acres of *L. latifolium* mapped, which increased to 245 acres in 2021, a 19.1% increase over three years (Figure 6). This is the largest percent change of any non-native species of concern in the Marsh since the 2018 remap. 223 acres occur in leveed areas (a 29% increase since 2015)

and 22 acres occur in tidal areas (a 31% decrease since 2018). In 1999, there were 646.4 acres of the *Lepidium* mapping unit, indicating a 62% decrease overall from 1999 to 2021 (Appendix G). Due to imagery limitations, not including alliances where *Lepidium* is only a component, and not indicating *L. latifolium* presence throughout the Marsh, the actual acreage of *Lepidium* is likely higher than these numbers indicate.


The non-native variety of *Phragmites australis* is the most widespread non-native species of concern in the marsh (Figure 6). This and the previous mapping efforts do not distinguish between the native and non-native forms; however, what is mapped as the *Phragmites australis* mapping unit is presumed to be the non-native form, which forms dense, monotypic stands. Since 1999, *P. australis* has increased by 540% over the entire marsh, from 693.1 acres in 1999 to 4,440 acres in 2021 (Appendix G). In 2015 there were 4,296 acres mapped, for an increase of 3.3% over three years. As of 2021, *P. australis* covered 2,530 acres (4.6%) of the leveed marsh (a 4% increase since 2018) and 1910.8 acres (17.1%) of the vegetation/land use tidal marsh (a 2.5% increase since 2018). While *Phragmites australis* continues to expand its cover, the rate of increase seems to have slowed compared to the period between 2015 and 2018, where *P. australis* had increased in cover by 45.8%.

While *Salsola soda* decreased in cover from 67 acres in 2015 to 39.1 acres in 2018, its cover has rebounded to 68 acres in 2021 (Figure 6). Our results indicate this increase could be due to differences in imagery potentially not fully capturing the discernable signature of this species in previous years.


**Non-native species abundance by management region:** All of the results of change detection for non-native species of concern are presented in Figures 7-13 as well as in Appendix G and are summarized or highlighted in the discussion below.




**Figure 7: *Arundo donax* distribution by management region between 1999, 2015, 2018, and 2021**




**Figure 8:** *Carpobrotus edulis* distribution by management region between 1999, 2015, 2018, and 2021




**Figure 9:** *Cortaderia selloana* distribution by management region between 1999, 2015, 2018, and 2021




**Figure 10: *Eucalyptus* spp. distribution by management region between 1999, 2015, 2018, and 2021**



**Figure 11: *Lepidium latifolium* distribution by management region between 1999, 2015, 2018, and 2021**



**Figure 12: *Phragmites australis* distribution by management region between 1999, 2015, 2018, and 2021**



**Figure 13: *Salsola soda* distribution by management region between 1999, 2015, 2018, and 2021**



**Region 0:**

Management Region 0 covers 25,024 acres (including 24,713 acres of Suisun Bay and sloughs). This region has a total of 310 acres of vegetation.

In 2021 this region included 0.37 acres of *Arundo donax*, which constitutes 4% of *Arundo donax* in the marsh (Figure 7).

No acres of *Carpobrotus edulis* were mapped in this region in previous years. In 2021 there was a 0.04-acre inclusion of *Carpobrotus edulis* in this region (Figure 8).

No acres of *Cortaderia selloana* were mapped in this region in previous years, however in 2021 there were 0.01 acres included in the region (Figure 9).

The cover of *Eucalyptus* spp. increased from 3.4 acres to 4 acres in 2021 (Figure 10). This is an overall 18% in cover over the 3-year period.

*Lepidium latifolium* decreased in this region from .09 acres in 2018 to .05 acres in 2021, indicating a 44% decrease in cover for this region (Figure 11).

There were 9 acres of *Phragmites australis* throughout the region in 2018, and the cover of *Phragmites* has since increased to 14 acres (Figure 12). This is a 56% increase over the 3-year period.

There was no *Salsola soda* detected in this region in 2021 or previous years.

**Region 1:**

Management Region 1 covers 18,988 acres (29% of the Marsh, excluding Suisun Bay and sloughs).

27% of the *Arundo donax*, 5.5% of the *Carpobrotus edulis*, 97% of the *Cortaderia selloana*, 32% of the *Eucalyptus* spp., 18% of the *Lepidium latifolium*, 22% of the *Phragmites australis*, and 38% of the *Salsola soda* occur within this management region.

*Arundo donax* has increased by 528% in this region since 1999. The increase from 2018 to 2021 has been 57%. There has been a 1.6-acre increase since 2018 (Figure 7).

*Carpobrotus edulis* has shown a 14% decrease from 2.5 acres in 2018 to 2.2 acres in 2021 (Figure 8). No *Carpobrotus* was mapped in this region in 1999.

*Cortaderia* has decreased by 31% since 2018, from 6.2 acres to 4.3 acres (Figure 9). There has been an overall 43% decrease since 1999 (Appendix G).

*Eucalyptus* spp. Has remained relatively static in this region since 2018. There has been a decrease of 1 acre from 103 acres in 2018 to 102 acres in 2021, a decrease of less than 1% (Figure 10). There has been an overall 94% increase since 1999 (Appendix G).

Vegetation dominated by *Lepidium latifolium* has decreased by 12% (6 acres) in this region since 2018 (Figure 11). There are currently 43 acres.

*Phragmites australis* has increased in this region by 794 acres since 1999 and 84 acres since 2018, for a total of 979 acres of *P. australis* in this region in 2021 (Figure 12). *P. australis* cover has increased by 9.4% over the 3-year period.

## 2021 Vegetation Map Update for Suisun Marsh

*Salsola soda* increased to 26 acres in 2021, an 89% increase since 2018 (Figure 13). All stands in this region, and throughout the Marsh in general, occur in leveed areas.

### **Region 2:**

Management Region 2 covers 9,958 acres (15% of the Marsh, excluding Suisun Bay and sloughs).

Five percent of the *Arundo donax*, 3% of the *Carpobrotus edulis*, 0.007% of the *Cortaderia selloana*, 9% of the *Eucalyptus* spp., 18% of the *Lepidium latifolium*, and 12% of the *Phragmites australis* occur within this management region. There were no mapped stands of *Salsola soda*.

There were 0.38 acres of *Arundo donax* in this region in 2021; a 24% decrease since there were 0.5 acres in 2018 (Figure 7).

*Carpobrotus edulis* has decreased from 1.4 acres in 2018 to 1.24 acres in 2018 (Figure 8).

There were 0.02 acres of *Cortaderia selloana* in 2018 and there has only been an increase of 0.01 acres for a total of 0.03 acres in 2021 (Figure 9).

There are 27 acres of *Eucalyptus* spp. in this region, a decrease of 5.5% since 2018 (Figure 10).

Vegetation dominated by *Lepidium latifolium* has decreased in this region by 3.8% (31 acres) since 2018 (Figure 11). There are currently 43 acres mapped in this region.

*Phragmites australis* has increased from 495 acres in 2018 to 551 acres in 2021, an 11% increase (Figure 12). Overall, there has been a 679% increase in this region since 1999.

*Salsola soda* was mapped in this region for the first time in 2015, with 8.2 acres in leveed areas. In 2021, there were no detectable stands present.

### **Region 3:**

Management Region 3 covers 4,720 acres (7% of the Marsh, excluding Suisun Bay and sloughs).

2% of the *Eucalyptus* spp. in the entire Marsh was found in this region, as was 2% of the *Lepidium latifolium*, and 5% of the *Phragmites australis*. There were no mapped stands of *Arundo donax*, *Carpobrotus edulis*, *Cortaderia selloana*, or *Salsola soda* in this region.

*Eucalyptus* spp. has shown no increase since 2018 and currently covers 7.5 acres (Figure 10).

There are 5.93 mapped acres of *Lepidium latifolium* in 2021, a 2.2% increase since 2021 (Figure 11).

*Phragmites australis* is the most abundant species of concern in this region. In the leveed areas, there are 94 acres, a 31% decrease since (Figure 12). There has been a 31% increase in tidal areas, from 101 acres in 2018 to 131 acres in 2021.

**Region 4:**

Management Region 4 covers 32,796 acres (49% of the Marsh, excluding Suisun Bay and sloughs).

45% of the *Arundo donax*, 91% of the *Carpobrotus edulis*, less than 1% of the *Cortaderia selloana*, 56% of the *Eucalyptus* spp., 62% of the *Lepidium latifolium*, 60% of the *Phragmites australis*, and 62% of the *Salsola soda* occur within this management region.

*Arundo donax* has decreased from 6.4 acres in 2018 to 4.15 acres in 2021, a 35% decrease for this region (Figure 7).

*Carpobrotus edulis* has increased from 36 acres in 2018 to 37 acres in 2021, a 1.6% increase for the region (Figure 8).

*Cortaderia selloana* cover dropped by 3.4 acres overall in this region, a decrease of 97% since 2018 (Figure 9).

*Eucalyptus* spp. has increased from 177 acres in 2018 to 179 acres in 2021 (Figure 10). 98% of stands occur in leveed areas.

*Lepidium latifolium* has increased by 44% in this region since 2018, from 106 acres to 153 acres (Figure 11). 97% of stands occur in leveed areas.

*Phragmites australis* is the most abundant species of concern in this region, covering a total of 2,670 acres. It has decreased in tidal areas by 4% (43 acres) since 2018. It has increased by 3% (54 acres) in the leveed areas of the region since 2018 (Figure 12). It has increased by 550% (2,260 acres) since 1999 (Appendix G).

*Salsola soda* was mapped at 25 acres in 2018 and has increased by 67% since 2018 to 42 acres (Figure 13). All stands occurred in leveed areas.

## Discussion, Conclusions, and Recommendations

### **Salt Marsh Harvest Mouse Habitat**

Total SMHM habitat has decreased since 2018 (a total decrease of 2,444 acres and 5% throughout the marsh) with a majority observed in leveed areas, where there was a 2,640-acre decrease, which reflects a 6.6% decrease in total leveed SMHM habitat (Appendix F). The amount of total tidal SMHM habitat increased by 196 acres and 2.23%.

While there has been an overall decrease in total SMHM habitat, the amount of habitat remains higher than 2015 levels, when there were 43,683 acres of SMHM habitat compared to 46,662 in 2021. However, total SMHM habitat has dipped below 1999 levels when there were 47,105.3 acres, and there has been an overall 0.94% decrease in total SMHM habitat since 1999.

Some of the observed decreases in SMHM habitat may be due to the refinement of road mapping for this mapping effort. Mappers were encouraged to extract road cover from vegetation polygons, and as a result, there was a 241-acre increase in road cover for this mapping effort compared to 2018. Roads often run through vegetation areas that are mapped with the Mediterranean California Naturalized Annual and Perennial Grassland Group mapping unit, which may account for a portion of the acreage decrease for this group.

Additionally, there was a substantial increase in acreage with the “Bare Ground” mapping unit (this unit is up over 2,000 acres since 2018). As illustrated in Figure 14 (see next page), following periods of flooding, areas that were once covered in vegetation can appear to be bare ground for short (or potentially long) periods. Mappers make attribution calls based on the imagery flown for the mapping effort, so the potential for vegetation to rebound in these areas is not considered. It may be that the habitat will be recovered once the habitat has been able to regrow post-perturbation, but mappers do not predict future growth when attributing.

**Figure 14: *Salicornia pacifica* pictured in 2018. The stand was flooded in 2020 and appears barren in 2021.**



**2018**



**2020**



**2021**

### Change in Tidal Habitat

For this mapping effort, mappers took note of levee breaches (both intentional and unintentional) that affect the habitat attribution of vegetation. Mappers found several examples of levee destruction that changed previously leveed habitat into tidal habitat. The newly formed nexus between sloughs and vegetation allows the habitat to be tidally influenced. Figure 15 below illustrates the demolition of a levee between 2018 and 2021 that has connected a slough to previously leveed habitat.

**Figure 15: A levee is demolished between 2018 and 2021.**



**2018**



**2020**



**2021**

### Non-Native Species of Concern

*Phragmites australis* is continuing to increase in cover; however, the rate of increase has reduced greatly since the 2018 remap. Since 2018, cover increased by 3.38%, whereas the rate of increase between 2015 and 2018 was 45%.

*Lepidium latifolium*, while at a much lower cover than *Phragmites*, has shown a greater rate of increase in recent years. From 2018 to 2021, *Lepidium* cover increased by 19%. Additionally, it may be that *Lepidium* is under-reported due to identification difficulty when it is not in flower at the time imagery is taken.

*Salsola soda* has made a rebound in its cover since 2018 and has returned to the levels it was recorded for the 2015 map. This can partially be attributed to the help of reconnaissance points that helped identify areas of rebound, and it should be noted that there was a 75% increase in cover for this species from 2018 to 2021.

*Arundo donax*, *Carpobrotus edulis*, *Cortaderia selloana*, and *Eucalyptus* spp. were all originally considered species of concern when mapping began. However, none of these species have shown increases in coverage (excluding increases due to better imagery) in recent years.

### Changes in Slough Cover

This mapping effort yielded a 489-acre decrease in slough cover. Previous maps of the bay from 2015 and 2018 misidentified a large water confluence for a slough, when in reality this confluence was not connected to the Suisun Bay and was surrounded by levees. Mappers correctly identified the habitat types for the vegetation surrounding this area, and the habitat types remained unaffected by this error. This error was corrected in the 2021 effort.

### Future Mapping Efforts

The Suisun map requires fine-scale vegetation classification and spatial delineation, which involves intensive effort on the part of ecologists. Consequently, the final map may be completed well after the base imagery is flown, a time lag that may not be acceptable to decision-makers. Vegetation mapping methods have changed with technology over the history of this project, becoming more efficient and accurate. However, we do not believe that fully automated mapping can achieve the same thematic accuracy as an experienced ecologist-mapper at this time. The time lag may be reduced by a combination of automated mapping techniques and manual attribution. Accuracy may be enhanced using a drone to acquire reconnaissance samples since access to field surveys is often hampered by flooded ditches or channels. A reevaluation of the required mapping resolution as well as periodic assessment of alternative mapping techniques, imagery acquisition, and field methods is recommended. Additionally, it may be beneficial for mappers to participate in data collection on the ground, to better understand ecological patterns of growth for the many species that occupy the marsh.

Data collected for this project included several surveys of species that are not included as map units for this project and are instead mapped at the group level. Future mapping efforts could include a review of



## 2021 Vegetation Map Update for Suisun Marsh

the data to identify consistent patterns of new species that could be added to the mapping hierarchy and evaluated as potential habitat. These additions should not be made based on a singular survey of a new species but rather repeated surveying of a new species throughout the marsh.

## Literature Cited

- Baldwin, B.G, D.H. Goldman, D.J. Keil, R. Patterson, T.J. Rosatti and D.H. Wilken, editors. 2012. The Jepson manual: vascular plants of California, second edition. University of California Press, Berkeley.
- [Boul, R. and T. Keeler-Wolf. 2008. 2006 Vegetation Map Update for Suisun Marsh, Solano County, California. A Report to the California Department of Water Resources. VegCAMP, California Department of Fish and Wildlife, Sacramento, CA.](#)
- [Boul, R. and T. Keeler-Wolf. 2016. 2012 Vegetation Map Update for Suisun Marsh, Solano County, California. A Report to the California Department of Water Resources. VegCAMP, California Department of Fish and Wildlife, Sacramento, CA.](#)
- California Department of Water Resources (DWR). 2013. Bay Delta Conservation Plan. Public Draft. November. Sacramento, CA. Prepared by ICF International (ICF 00343.12). Sacramento, CA.
- Congalton, R. and K. Green. 1999. Assessing the accuracy of remotely sensed data: principles and practices. Lewis Publishers, New York.
- Federal Geographic Data Committee (FGDC). 2008. National Vegetation Classification Standard, Version 2 FGDC-STD-005-2008 (version 2). Vegetation Subcommittee, FGDC Secretariat, U.S. Geological Survey. Reston, VA. 55 pp. + Appendices.
- Foody, G. 2002. Status of land cover classification accuracy assessment. Remote Sens Environ 80:185–201.
- [Geographical Information Center \(GIC\). 2022. 2018 Vegetation Map Update for Suisun Marsh Solano County, California. A Report to the California Department of Water Resources. Geographical Information Center, North State Planning and Development Collective, California State University, Chico, CA.](#)
- Gopal, S. and C. Woodcock. 1994. Theory and methods for accuracy assessment of thematic maps using fuzzy sets. Photogramm Eng Remote Sens 60:181–188.
- Hagen, A. 2003. Fuzzy set approach to assessing similarity of categorical maps. Int J Geogr Inf Sci 17(3):235–249.
- [Keeler-Wolf, T. and M. Vaghti. 2000. Vegetation Mapping of Suisun Marsh, Solano County – A Report to the California Department of Water Resources. Unpublished administrative report on file at Biogeographic Data Branch, California Department of Fish and Wildlife, Sacramento.](#)
- Kincaid, T. and A. Olsen. 2011. survey: Spatial Survey Design and Analysis. R package version 2.2.
- Lea, C. and A. Curtis. 2010. Thematic accuracy assessment procedures: National Park Service Vegetation Inventory, version 2.0. Natural Resource Report NPS/2010/NRR—2010/204. National Park Service, Fort Collins, Colorado.

## 2021 Vegetation Map Update for Suisun Marsh

Metzler, J. and S. Sader. 2005. Agreement assessment of spatially explicit regression-derived forest cover and traditional forest industry stand type maps. *Photogramm Eng Remote Sens* 71(11):1303–1309.

San Francisco Estuary Institute (SFEI). 2011. Bay Area Aquatic Resources Inventory (BAARI) Basemap.

Story, M. and R. Congalton. 1986. Accuracy Assessment: A User's Perspective. *Photogrammetric Engineering and Remote Sensing* 52:397–399.

Sustaita, D., P. Frinfrock Quickert, L. Patterson, L. Barthman-Thompson, S. Estrella. Salt marsh harvest mouse demography and habitat use in the Suisun Marsh, California. *Journal of Wildlife Management* 75(6):1498-1507.

USDA NRCS. 2015. The PLANTS Database (<http://plants.usda.gov>, 4 June 2015). National Plant Data Team, US Department of Agriculture, Natural Resources Conservation Service, Greensboro, NC 27401-4901 USA.

[USDA Bureau of Reclamation, U.S. Fish and Wildlife Service, and California Department of Fish and Game. 2013. Suisun Marsh Habitat Management, Preservation and Restoration Plan. May 2013.](#)

[Vaghti, M., T. Keeler-Wolf. 2001. Suisun Marsh Vegetation Mapping Change Detection 2000. A Report to the California Department of Water Resources. Wildlife and Habitat Data Analysis Branch, California Department of Fish and Wildlife, Sacramento.](#)

[Vaghti, M. and T. Keeler-Wolf. 2004. Suisun Marsh Vegetation Mapping Change Detection 2003. A Report to the California Department of Water Resources. Wildlife and Habitat Data Analysis Branch, California Department of Fish and Wildlife, Sacramento.](#)

[Vegetation Classification and Mapping Program \(VegCAMP\). 2012. 2009 Vegetation map update for Suisun Marsh, Solano County, California. A report to the California Department of Water Resources. VegCAMP, California Department of Fish and Wildlife, Sacramento, CA.](#)

[Vegetation Classification and Mapping Program \(VegCAMP\). 2018. 2015 Vegetation map update for Suisun Marsh, Solano County, California. A report to the California Department of Water Resources. VegCAMP, California Department of Fish and Wildlife, Sacramento, CA.](#)

## Appendix A

### Suisun Marsh Vegetation Classification and Hierarchy

The vegetation and mapping classification hierarchy of Suisun Marsh is compliant with the United States National Vegetation Classification (USNVC). The vegetation types are listed below within the full eight levels of the USNVC hierarchy. Each type that has been used in any Suisun Marsh map is followed by its mapping code in parentheses.

The original vegetation classification for Suisun Marsh was created in 1999. Since then, a great deal of vegetation work has been done in California, and the state and national hierarchies have been refined. Following the 2015 remap and accuracy assessment, the Suisun Marsh vegetation types were reviewed and modified to match the most current USNVC hierarchy more closely. Some types that were difficult to distinguish on the imagery were aggregated to higher levels of the hierarchy (group or macrogroup level). Some types that were equivalent to current USNVC types were renamed to conform to the current hierarchy. New mapping codes have been assigned to all vegetation types as of the 2015 map; the new codes are listed in parentheses following each type.

Although many alliances and associations from the original classification are equivalent to types in the current USNVC hierarchy, some have no comparable current types. These unmatched vegetation types are labeled “mapping units” and are placed within the appropriate alliances and groups in the hierarchy structure below.

Some of the original mapping units were defined too broadly to fit into the alliance and association levels of the current hierarchy. However, they do approximate some of the upper levels. For instance, although the Medium Upland Graminoids Mapping Unit does not fit into the species-oriented alliance structure, it can be considered part of the Mediterranean California Naturalized Annual and Perennial Grassland Group. These broadly defined mapping units are listed with their corresponding USNVC vegetation types under “Legacy Mapping Units” at the end of the hierarchy.

In the future, when the Suisun Marsh vegetation field data are reanalyzed with a larger data set, the mapping units will either be confirmed as distinct types or renamed to better fit an existing type. In future maps, the broadly defined mapping units will not be used but will be replaced by the corresponding USNVC vegetation types.

The USNVC hierarchy is composed of eight levels, organized into three upper, three middle, and two lower levels as shown below:

<u>Level</u>	<u>Example</u>
Upper	-
Level 1 - Class	Forest and Woodland
Level 2 - Subclass	Temperate Forest
Level 3 - Formation	Warm Temperate Forest
Middle	-
Level 4 - Division	Madrean Forest and Woodland
Level 5 - Macrogroup	California Forest and Woodland
Level 6 - Group	Californian Broadleaf Forest and Woodland
Lower	-
Level 7 - Alliance	<i>Quercus agrifolia</i>

Descriptions of vegetation types in the 2015 map can be found in Appendix B.

**1. Forest and Woodland Class**

**a. Temperate Forest Subclass**

**i. Warm Temperate Forest Formation (1000000)**

**1. Madrean Forest and Woodland Division (1100000)**

**a. California Forest and Woodland Macrogroup (1110000)**

**i. Californian Broadleaf Forest and Woodland Group<sup>1</sup> (1111000)**

**1. *Quercus agrifolia* Woodland Alliance (1111100)**

**a. *Quercus agrifolia* Association (1111101)**

**b. *Quercus agrifolia* / *Salix lasiolepis* Association (1111102)**

**2. *Quercus lobata* Woodland Alliance (1111200)**

**ii. Cool Temperate Forest Formation (2000000)**

**1. North American Introduced Evergreen Broadleaf and Conifer Forest Division (2100000)**

**a. Introduced North American Mediterranean Woodland and Forest Macrogroup (2110000)**

**i. no subdivision at group level**

**1. *Eucalyptus (globulus, camaldulensis)* Woodland Semi-Natural Alliance (2110100)**

**a. *Eucalyptus globulus* Semi-Natural Association (2110101)**

**2. *Ailanthus altissima* Woodland Semi-Natural Alliance (2110200)**

**b. Southwestern North American Riparian, Flooded and Swamp Forest Macrogroup (2120000)**

**i. Southwestern North American Riparian Evergreen and Deciduous Woodland Group<sup>2</sup> (2121000)**

**1. *Salix laevigata* Woodland Alliance (2121100)**

**a. *Salix laevigata* / *Salix lasiolepis* Association (2121101)**

**ii. Southwestern North American Riparian/Wash Scrub Group (2122000)**

**1. *Salix exigua* Shrubland Alliance (2122100)**

**a. *Salix exigua* Association (2122101)**

**2. *Rosa californica* Shrubland Alliance (2122200)**

**a. *Rosa californica* Association (2122201)**

**b. *Rosa californica* – *Baccharis pilularis* Association (2122202)**

**3. *Baccharis salicifolia* Shrubland Alliance (2122300)**

**iii. Southwestern North American Introduced Riparian Scrub Group (2123000)**

**1. *Arundo donax* Herbaceous Semi-Natural Alliance (2123100)**

**a. *Arundo donax* Semi-Natural Association (2123101)**

**2. *Tamarix* spp. Shrubland Semi-Natural Alliance (2123200)**

**iii. Temperate Flooded and Swamp Forest Formation (3000000)**

**1. Western North America Flooded and Swamp Forest Division (3100000)**

**a. Western Cordilleran Montane–Boreal Riparian Scrub Macrogroup (3110000)**

---

<sup>1</sup> Formerly named Oaks Mapping Unit

<sup>2</sup> Formerly named Willow Trees Mapping Unit

- i. Vancouverian Riparian Deciduous Forest Group (3111000)
  - 1. *Fraxinus latifolia* Woodland Alliance (3111100)
    - a. *Fraxinus latifolia* Planted Stands Mapping Unit (3111101)
- 2. Temperate Flooded and Swamp Forest Mesomorphic Shrub and Herb Vegetation Class**
  - a. Mediterranean Scrub and Grassland Subclass**
    - i. Mediterranean Grassland and Forb Meadow Formation (4000000)**
      - 1. California Grassland and Meadow Division (4100000)**
        - a. California Annual and Perennial Grassland Macrogroup (4110000)**
          - i. Mediterranean California Naturalized Annual and Perennial Grassland Group<sup>3</sup> (4111000)
            - 1. *Brassica nigra* and Other Mustards Herbaceous Semi-Natural Alliance (4111100)
              - a. *Brassica nigra* Semi-Natural Association (4111101)
              - b. *Raphanus sativus* Semi-Natural Association (4111102)
            - 2. *Bromus (diandrus, hordeaceus) – Brachypodium distachyon* Herbaceous Semi-Natural Alliance (4111200)
            - 3. *Centaurea (solstitialis, melitensis)* Herbaceous Semi-Natural Alliance (4111300)
            - 4. *Conium maculatum – Foeniculum vulgare* Herbaceous Semi-Natural Alliance (4111400)
              - a. *Conium maculatum* Semi-Natural Association (4111401)
              - b. *Foeniculum vulgare* Semi-Natural Association (4111402)
            - 5. *Cortaderia (jubata, selloana)* Herbaceous Semi-Natural Alliance (4111500)
            - 6. *Festuca perennis* Herbaceous Semi-Natural Alliance (4111600)
              - a. *Festuca perennis – Lepidium latifolium* Semi-Natural Association (4111601)
              - b. *Festuca perennis – Lotus corniculatus* Semi-Natural Association (4111602)
              - c. *Festuca perennis – Rumex* spp. Mapping Unit (4111603)
              - d. *Hordeum marinum – Festuca perennis* Mapping Unit (4111604)
            - 7. *Elytrigia pontica* Mapping Unit (4111700)
            - 8. *Agrostis avenacea* Mapping Unit (4111800)
            - 9. *Vulpia* spp. – *Euthamia occidentalis* Mapping Unit (4111900)
  - 3. Mesomorphic Shrub and Herb Vegetation Class**
    - a. Temperate and Boreal Shrubland and Grassland Subclass**
      - i. Temperate Grassland, Meadow, and Shrubland Formation (5000000)**
        - 1. Vancouverian and Rocky Mountain Grassland and Shrubland Division (5100000)**
          - a. Western North American Temperate Grassland and Meadow Macrogroup (5110000)**
            - i. Vancouverian and Rocky Mountain Naturalized Perennial Grassland Group (5111000)
              - 1. *Phalaris aquatica* Herbaceous Semi-Natural Alliance (5111100)
                - a. *Phalaris aquatica* Semi-Natural Association (5111101)
        - b. Vancouverian Lowland Grassland and Shrubland Macrogroup (5120000)**
          - i. Naturalized Non-Native Deciduous Scrub Group (5121000)
            - 1. *Rubus armeniacus* Shrubland Semi-Natural Alliance (5121100)
      - ii. Temperate and Boreal Scrub and Herb Coastal Vegetation Formation (6000000)**

---

<sup>3</sup> Formerly named Annual Grasses/Weeds Mapping Unit

- 1. Pacific Coast Scrub and Herb Littoral Vegetation Division (6100000)**
  - a. Vancouverian Coastal Dune and Bluff Macrogroup (6110000)**
    - i. California Coastal Evergreen Bluff and Dune Scrub Group<sup>4</sup> (6111000)
      1. *Baccharis pilularis* Shrubland Alliance (6111100)
        - a. *Baccharis pilularis* / Annual Grass–Herb Association (6111101)
    - ii. California–Vancouverian Semi-Natural Littoral Scrub and Herb Vegetation Group (6112000)
      1. *Carpobrotus edulis* or Other Ice Plants Herbaceous Semi-Natural Alliance (6112100)
- iii. Temperate and Boreal Freshwater Marsh Formation (7000000)**
  - 1. Western North American Freshwater Marsh Division (7100000)**
    - a. Western North American Freshwater Marsh Macrogroup (7110000)**
      - i. Arid West Freshwater Emergent Marsh Group<sup>5</sup> (7111000)
        1. *Typha (angustifolia, latifolia, domingensis)* Herbaceous Alliance (7111100)
          - a. *Typha (angustifolia, latifolia, domingensis)* Association (7111101)
          - b. *Typha (angustifolia, latifolia, domingensis)* (dead stalks) Mapping Unit (7111102)
          - c. *Typha (angustifolia, latifolia, domingensis)* – *Echinochloa crus-galli* Association (7111103)
          - d. *Typha (angustifolia, latifolia, domingensis)* – *Distichlis spicata* Association (7111104)
          - e. *Typha (angustifolia, latifolia, domingensis)* – *Phragmites australis* Association (7111105)
          - f. *Typha (angustifolia, latifolia, domingensis)* – *Schoenoplectus americanus* Association (7111106)
        2. *Schoenoplectus (acutus, californicus)* Herbaceous Alliance (7111200)
          - a. *Schoenoplectus californicus* – *Schoenoplectus acutus* Association (7111201)
          - b. *Schoenoplectus (acutus, californicus)* Mapping Unit (7111202)
          - c. *Schoenoplectus (acutus, californicus)* – *Rosa californica* Association (7111203)
          - d. *Schoenoplectus (acutus, californicus)* – Wetland Herbs Mapping Unit (7111204)
        3. *Schoenoplectus (acutus, californicus)* – *Typha (angustifolia, latifolia, domingensis)* Mapping Unit (7111300)
        4. *Phragmites australis* Herbaceous Alliance (7111400)
          - a. *Phragmites australis* Association (7111401)
          - b. *Phragmites australis* – *Schoenoplectus* spp. Association (7111402)
          - c. *Phragmites australis* – *Xanthium strumarium* Mapping Unit (7111403)
        5. *Alisma triviale* Mapping Unit (7111500)
      - ii. Vancouverian Coastal/Tidal Marsh and Meadow Group (7112000)
        1. *Potentilla anserina* Herbaceous Alliance (7112100)

---

<sup>4</sup> Formerly named Medium Upland Shrubs Mapping Unit

<sup>5</sup> Formerly named Tall Wetland Graminoids Mapping Unit



**b. Western North America Vernal Pool Macrogroup (7120000)**

- i. California Vernal Pool and Grassland Matrix Group (7121000)
- ii. Californian Mixed Annual/Perennial Freshwater Vernal Pool/Swale Bottomland Group (7121000)
  - 1. *Grindelia (stricta)* Provisional Herbaceous Alliance (7121100)

**c. Western North America Wet Meadow and Low Shrub Carr Macrogroup<sup>6</sup> (7130000)**

- i. Californian Warm Temperate Marsh/Seep Group (7131000)
  - 1. *Juncus arcticus* (var. *balticus*, *mexicanus*) Herbaceous Alliance (7131100)
    - a. *Juncus arcticus* var. *balticus* Association (7131101)
    - b. *Juncus arcticus* var. *balticus* – *Conium maculatum* Association (7131102)
    - c. *Juncus arcticus* var. *balticus* – *Lepidium latifolium* Association (7131103)
    - d. *Juncus arcticus* var. *balticus* – *Potentilla anserina* Association (7131104)
  - 2. *Leymus triticoides* Herbaceous Alliance (7131200)
- ii. Naturalized Warm-Temperate Riparian And Wetland Group (7132000)
  - 1. *Lepidium latifolium* Herbaceous Semi-Natural Alliance (7132100)
  - 2. *Persicaria lapathifolia* – *Xanthium strumarium* Herbaceous Alliance (7132200)
    - a. *Persicaria* spp. – *Xanthium strumarium* – *Echinochloa crus-galli* Mapping Unit (7132201)
  - 3. *Cynodon dactylon* – *Crypsis* spp. – *Paspalum* spp. Moist Herbaceous Semi-Natural Alliance (7132300)
    - a. *Crypsis schoenoides* Mapping Unit (7132301)
    - b. *Cynodon dactylon* Mapping Unit (7132302)
  - 4. *Polypogon monspeliensis* Mapping Unit (7132400)
  - 5. *Rumex* spp. Mapping Unit (7132500)
  - 6. *Salsola soda* Mapping Unit (7132600)

**iv. Temperate and Boreal Salt Marsh Formation (8000000)**

**1. Temperate and Boreal Pacific Coastal Salt Marsh Division (8100000)**

**a. North American Pacific Coastal Salt Marsh Macrogroup<sup>7</sup> (8110000)**

- i. Temperate Pacific Tidal Salt and Brackish Meadow Group (8111000)
  - 1. *Bolboschoenus maritimus* Herbaceous Alliance (8111100)
    - a. *Bolboschoenus maritimus* Association (8111101)
    - b. *Bolboschoenus maritimus* – *Salicornia pacifica* Association (8111102)
    - c. *Bolboschoenus maritimus* – *Sesuvium verrucosum* Association (8111103)
  - 2. *Distichlis spicata* Herbaceous Alliance (8111200)
    - a. *Distichlis spicata* Association (8111201)
    - b. *Distichlis spicata* – Annual Grasses Association (8111202)
    - c. *Distichlis spicata* – *Juncus arcticus* var. *balticus* (*J. arcticus* var. *mexicanus*) Association (8111203)
    - d. *Distichlis spicata* – *Lotus corniculatus* Mapping Unit (8111204)
    - e. *Distichlis spicata* – *Salicornia pacifica* Association (8111205)
    - f. *Distichlis spicata* – *Schoenoplectus americanus* Provisional Association (8111206)

---

<sup>6</sup> Formerly named Medium Wetland Graminoids Mapping Unit

<sup>7</sup> Formerly named Short Wetland Graminoids Mapping Unit

- g. *Distichlis spicata* – *Cotula coronopifolia* Association (8111207)
- h. *Distichlis spicata* – *Bolboschoenus maritimus* Mapping Unit (8111208)
- i. *Distichlis spicata* – *Juncus balticus* – *Triglochin* spp. – *Glaux maritima* Mapping Unit (8111209)
- j. *Lepidium latifolium* – *Distichlis spicata* Semi-Natural Association (8111210)
- 3. *Salicornia pacifica* (*Salicornia depressa*) Herbaceous Alliance (8111300)
  - a. *Salicornia pacifica* Association (8111301)
  - b. *Salicornia pacifica* – Annual Grasses Association (8111302)
  - c. *Salicornia pacifica* – *Atriplex prostrata* Association (8111303)
  - d. *Salicornia pacifica* – *Crypsis schoenoides* Association (8111304)
  - e. *Salicornia pacifica* – *Sesuvium verrucosum* Association (8111305)
  - f. *Salicornia pacifica* – *Echinochloa crus-galli* – *Polygonum* – *Xanthium strumarium* Association (8111306)
  - g. *Salicornia pacifica* – *Cotula coronopifolia* Association (8111307)
- 4. *Spartina foliosa* Herbaceous Alliance (8111400)
- ii. Western North American Disturbed Alkaline Marsh and Meadow Group (8112000)
  - 1. *Atriplex prostrata* – *Cotula coronopifolia* Herbaceous Semi-Natural Alliance (8112100)
    - a. *Cotula coronopifolia* Semi-Natural Association (8112101)
    - b. *Atriplex prostrata* Semi-Natural Association (8112102)
    - c. *Atriplex prostrata* – *Distichlis spicata* Semi-Natural Association (8112103)
    - d. *Atriplex prostrata* – *Bolboschoenus maritimus* Semi-Natural Association (8112104)
    - e. *Atriplex prostrata* – *Sesuvium verrucosum* Semi-Natural Association (8112105)
    - f. *Atriplex prostrata* – Annual Grasses Semi-Natural Association (8112106)
  - 2. *Sesuvium verrucosum* Herbaceous Alliance (8112200)
    - a. *Sesuvium verrucosum* Association (8112201)
    - b. *Sesuvium verrucosum* – *Distichlis spicata* Association (8112202)
    - c. *Sesuvium verrucosum* – *Festuca perennis* Association (8112203)
    - d. *Sesuvium verrucosum* – *Cotula coronopifolia* Association (8112204)
  - 3. *Spergularia marina* Provisional Herbaceous Alliance (8112300)
    - a. *Spergularia marina* – *Cotula coronopifolia* Mapping Unit (8112301)
- 2. Western North American Interior Alkali–Saline Wetland Division (8200000)**
  - a. Warm Semi-Desert/Mediterranean Alkali–Saline Wetland Macrogroup (8210000)**
    - i. Southwestern North American Alkali Marsh/Seep Vegetation Group (8211000)
      - 1. *Schoenoplectus americanus* Herbaceous Alliance (8211100)
        - a. *Schoenoplectus americanus* Association (8211101)
        - b. *Schoenoplectus americanus* – *Potentilla anserina* Association (8211102)
        - c. *Schoenoplectus americanus* – *Schoenoplectus californicus* – *Schoenoplectus acutus* Association (8211103)
        - d. *Schoenoplectus americanus* – *Lepidium latifolium* Association (8211104)

- ii. Southwestern North American Salt Basin and High Marsh Group (8212000)
  - 1. *Atriplex lentiformis* Shrubland Alliance (8212100)
  - 2. *Frankenia salina* Herbaceous Alliance (8212200)
    - a. *Frankenia salina* – Annual grasses Mapping Unit (8212201)
    - b. *Frankenia salina* – *Distichlis spicata* Association (8212202)
- 4. Hydromorphic Vegetation (Aquatic Vegetation) Class (9000000)**
  - a. Saltwater Aquatic Vegetation Subclass**
    - i. Marine and Estuarine Saltwater Aquatic Vegetation Formation (9100000)**
      - 1. Temperate Pacific Saltwater Aquatic Vegetation Division (9110000)**
        - a. Temperate Pacific Intertidal Shore Macrogroup (9111000)**
          - i. Temperate Pacific Intertidal Flat Group (9111100)
            - 1. *Stuckenia (pectinata)* – *Potamogeton* spp. Herbaceous Alliance (9111200)
              - a. *Stuckenia pectinata* Association (9111201)
            - 2. *Ruppia (cirrhosa, maritima)* Herbaceous Alliance (9111300)
  - b. Freshwater Aquatic Vegetation Subclass**
    - i. Freshwater Aquatic Vegetation Formation**
      - 1. North American Freshwater Aquatic Vegetation Division (9200000)**
        - a. Western North American Freshwater Aquatic Vegetation Macrogroup (9210000)**
          - i. Naturalized Temperate Pacific Freshwater Vegetation Group (9211000)
            - 1. *Ludwigia (hexapetala, peploides)* Provisional Herbaceous Semi-Natural Alliance (9211100)
- 5. Agriculture & Developed Vegetation Class**
  - a. Herbaceous Agricultural Vegetation Subclass (9310000)**
    - i. Row & Close Grain Crop Formation<sup>8</sup> (9311000)**
    - ii. Fallow Field and Weed Vegetation Formation (9312000)**
      - 1. Cropland Fallow Field Division (9312100)**
        - a. Fallow Field Macrogroup (9312110)**
  - b. Herbaceous & Woody Developed Vegetation Subclass (9410000)**
    - i. Other Developed Vegetation Formation (9411000)**
      - 1. Other Developed Vegetation Division (9411100)**
        - a. Tree Developed Vegetation Macrogroup (9411110)**
          - i. Temperate Tree Developed Vegetation Group<sup>9</sup> (9411111)

#### **Non-Vegetation Mapping Units**

- Bare Ground Mapping Unit (1)
- Agriculture (2)
- Road / Trails Mapping Unit (4)
- Slough Mapping Unit (6)
- Tidal Mudflat Mapping Unit (7)
- Railroad Track Mapping Unit (8)
- Ditch Mapping Unit (9)

---

<sup>8</sup> Formerly named Cultivated Annual Graminoid Mapping Unit

<sup>9</sup> Formerly named Landscape Trees Mapping Unit

## 2021 Suisun Marsh Vegetation Map Update for

Open Water Mapping Unit (11)

Developed Mapping Unit (15)

### **Legacy Mapping Units**

#### **Californian Broadleaf Forest and Woodland Group (1111000)**

Oaks Mapping Unit (1111001)

#### **Southwestern North American Riparian Evergreen and Deciduous Woodland Group (2121000)**

Willow Trees Mapping Unit (2121001)

#### **Southwestern North American Riparian/Wash Scrub Group (2122000)**

Tall Wetland Shrubs Mapping Unit (2122001)

Medium Wetland Shrubs Mapping Unit (2122002)

#### **California Annual and Perennial Grassland Macrogroup (4110000)**

Annual Grasses Mapping Unit (4110001)

#### **Mediterranean California Naturalized Annual and Perennial Grassland Group (4111000)**

Tall Upland Graminoids Mapping Unit (4111001)

Medium Upland Graminoids Mapping Unit (4111002)

Perennial Grass Mapping Unit (4111003)

Short Upland Graminoids Mapping Unit (4111004)

Tall Upland Herbs Mapping Unit (4111005)

Medium Upland Herbs Mapping Unit (4111006)

Short Upland Herbs Mapping Unit (4111007)

Annual Grasses/Weeds Mapping Unit (4111008)

#### **California Coastal Evergreen Bluff and Dune Scrub Group (6111000)**

Medium Upland Shrubs Mapping Unit (6111001)

#### **Arid West Freshwater Emergent Marsh Group (7111000)**

Tall Wetland Graminoids Mapping Unit (7111001)

*Calystegia sepium* – *Euthamia occidentalis* Mapping Unit (7111002)

#### **Western North America Wet Meadow and Low Shrub Carr Macrogroup (7130000)**

Medium Wetland Graminoids Mapping Unit (7130001)

#### **Temperate and Boreal Salt Marsh Formation (8000000)**

Wetland Herbs Mapping Unit (8000001)

Tall Wetland Herbs Mapping Unit (8000002)

Medium Wetland Herbs Mapping Unit (8000003)

#### **North American Pacific Coastal Salt Marsh Macrogroup (8110000)**

Short Wetland Graminoids Mapping Unit (8110001)

Short Wetland Herbs Mapping Unit (8110002)

#### **Hydromorphic Vegetation (Aquatic Vegetation) Class (9000000)**

Floating-leaved Wetland Herbs Mapping Unit (9000001)

#### **Row & Close Grain Crop Formation (9311000)**

Cultivated Annual Graminoid Mapping Unit (9311001)

#### **Temperate Tree Developed Vegetation Group (9411111)**

Landscape Trees Mapping Unit (9411112)

#### **Developed (15)**

Parking Lot Mapping Unit (3)

Structure Mapping Unit (5)

Water Treatment Pond Mapping Unit (13)

## 2021 Suisun Marsh Vegetation Map Update for

Urban Area Mapping Unit (14)

**Road / Trails Mapping Unit (4)**

Trail Mapping Unit (10)

## Appendix B

### Suisun Marsh Vegetation Type Descriptions

These descriptions were created for vegetation types mapped in Suisun Marsh in 2015.

These descriptions are intended for use as a guide to the identification of field-based and image interpretation-based vegetation assessments. An example of the type as it is represented in imagery is included with each description. Terms used in the descriptions are defined as follows:

**Alliance:** Plant communities based on dominant/diagnostic species of the uppermost or dominant stratum. Accepted alliances are part of the USNVC hierarchy.

**Association:** The most botanically detailed or finest-scale plant community designation based on dominant species and multiple co- or sub-dominant indicator species from any strata. Associations are part of the USNVC hierarchy.

**Plant community nomenclature:** Species separated by "-" are within the same stratum; species separated by "/" are in different strata. The number that follows some plant community names is the Mapping Code used for labeling plant community polygons for the associated GIS-based plant community map.

**Cover:** The primary metric used to quantify the importance/abundance of a particular species or a particular vegetation layer within a stand. It is measured by estimating the aerial extent of the living plants, or the bird's-eye view looking from above, for each category. Cover in this mapping project uses the concept of "porosity" or foliar cover rather than "opacity" or crown cover. Thus, field crews are trained to estimate the amount of light versus shade produced by the canopy of a plant or a stratum by considering the amount of shade it casts excluding the openings it may have in the interstitial spaces (e.g., between leaves or branches). This is assumed to provide a more realistic estimate of the actual amount of shade cast by the individual or stratum which, in turn, relates to the actual amount of light available to individual species or strata beneath it. However, as a result, cover estimates can vary substantially between leaf-on versus leaf-off conditions.

**Absolute cover:** The actual percentage of the surface area of the survey that is covered by a species or physiognomic group (trees, shrubs, herbaceous), as in "*Salicornia pacifica* covers 10% of the area being surveyed." Absolute cover of all species or physiognomic groups, when added together, may total greater than 100%, because this is not a proportional number, and plants can overlap each other. For example, a stand could have 25% tree cover in the upper layer, 40% shrub cover in the middle layer, and 50% herbaceous cover on the ground. However, when aerial interpretation is being used, the maximum absolute value is 100% since lower levels of vegetation cannot be seen through the overstory in aerial photographs.

**Relative cover:** The percentage of surface area within a survey area that is covered either by one species relative to other species within the same physiognomic stratum (tree, shrub, herbaceous) or one stratum relative to the total vegetation cover in a polygon. Thus, 50% relative cover of *Distichlis spicata* in the herb layer means that *D. spicata* comprises half the cover of all herbaceous species within a stand, while 50% relative shrub cover means that shrubs make up half the cover of all vegetation within a stand. Relative cover values are proportional numbers that, when added together, total 100% for all the species within a stratum or each stratum within a stand of vegetation.

**Dominance:** Dominance refers to the preponderance of vegetation cover in a stand of uniform composition and site history. It may refer to the cover of an individual species as in "dominated by *Salicornia pacifica*," or it may refer to dominance by a physiognomic group, as in "dominated by herbs." When we use the term in the key, a species is dominant if it is in relatively high cover in each stand, however, see "dominance by layer," below.

**Strongly dominant:** A species in the dominant life form stratum has 60% or greater relative cover.

**Co-dominant:** Co-dominance refers to two or more species in a stand with similar cover. Specifically, each species has between 30% to 60% relative cover. For example, in a coastal scrub stand with 5% *Baccharis pilularis*, 4% *Frangula californica*, and 3% *Rubus ursinus* (total 13% shrub cover), technically only the *Baccharis* ( $5/13 = 39\%$  relative cover) and the *Frangula* ( $4/13 = 31\%$  relative cover) would be co-dominant because *Rubus* would only have 23% relative cover ( $3/13 = 23\%$ ).

**Consistent/Characteristic/Diagnostic species:** Should be present in at least 80% of the stands of the type, with no restriction on cover. Relatively even spacing throughout the stand is important particularly in vegetation with low total cover since an even distribution of the diagnostic species is a much better indicator than overall cover. Characteristic species that are evenly distributed are better indicators of a type than species with higher cover and patchy distribution.

**Dominance by layer:** Tree, shrub, and herbaceous layers are considered physiognomically distinct. A vegetation type is considered to belong to a certain physiognomic group if it is dominated by one layer. Layers are prioritized in order of height. If the tallest layer is dominant and characteristic (see definitions above) across multiple stands of one type, the alliance is usually named by the dominant and/or characteristic species of the tallest layer. Average covers within the dominant layer reflect the "modal" concept of the health/age/environment of a particular vegetation type. For example, a higher average cover of woody plants within a stand not recently affected by disturbance reflects a mode of general availability of water, nutrition, and equitable climate, while lower average cover under similar conditions would reflect lower availability of these things.

**Woody plant:** A vascular plant species that has a noticeably woody stem (e.g., shrubs and trees). It does not include herbaceous species with woody underground portions such as tubers, roots, or rhizomes.

**Tree:** A one-stemmed woody plant that normally grows to be greater than 5 meters tall. In some cases, trees may be multi-stemmed (ramified due to fire or other disturbance) but the height of mature plants typically exceeds 5 meters. If less than 5 meters tall, undisturbed individuals of these species are usually single-stemmed. Certain species that can resemble either shrubs or trees (e.g., *Aesculus californica*) are, out of statewide tradition or by the USNVC, called trees. It behooves one to memorize which species are "traditionally" placed in one life form or another. We use the accepted lifeforms in the USNVC or the PLANTS Database (USDA NRCS 2015) to do this.

**Shrub:** A multi-stemmed woody plant that is usually 0.2–5 meters tall. Definitions are blurred at the low and high ends of the height scales. At the tall end, shrubs may approach tree-size based on disturbance frequencies (e.g., old-growth re-sprouting chaparral species such as *Cercocarpus montanus*, *Fremontodendron californicum*, *Prunus ilicifolia*, and so forth, may frequently attain "tree size," but are still typically multi-stemmed and are considered shrubs in this key). At the short end, woody perennial herbs or sub-shrubs of various species are often difficult to categorize into a consistent life form (e.g., *Eriogonum latifolium*, *Lupinus chamissonis*); in such instances, we refer to the PLANTS Database or "pick a lane" based on best available definitions.

*Subshrub*: A multi-stemmed plant with noticeably woody stems less than 0.5 meters tall. May be easily confused with a perennial herb or small shrub. We lump them into the “shrub” category in stand tables and descriptions of vegetation types.


*Shrub-characterized vegetation*: Shrubs (including sub-shrubs) are evenly distributed throughout the stand, providing a consistent (even if sparse) structural component; the stand cannot be characterized as a tree stand; and one or both of the following criteria are met: (1) shrubs influence the distribution or population dynamics of other plant species; (2) shrubs play an important role in ecological processes within the stand. For this project, shrub alliances have at least 10% absolute shrub cover.

*Herbaceous plant*: Any species of plant that has no main woody stem development; including grasses, forbs, and perennial species that die back each year.

*Herb-characterized vegetation*: Herbs are evenly distributed throughout the stand, providing a consistent (even if sparse) structural component and playing an important role in ecological processes within the stand. The stand cannot be characterized as a tree or shrub stand.

*Botanical nomenclature*: We use the PLANTS Database in vegetation mapping as our standard for botanical names. In certain cases, if a plant is recognized by *The Jepson Manual, second edition* (Baldwin et al. 2012), but not PLANTS Database, we default to the Jepson name.


### **Bare Ground Mapping Unit**



Ground generally has <10% cover of vegetation. Does not include unvegetated tidal flats but does include unvegetated leveed ground such as scraped areas or formerly inundated areas with no vegetation.



**Road / Trails Mapping Unit**



Includes roads and trails that are wider than the minimum mapping width of 10 feet. May have mown vegetation.

**Slough Mapping Unit**



Wide, fully tidal waterways.

**Tidal Mudflat Mapping Unit**




Tidally exposed mud deposits, generally unvegetated or with a low cover of herbs or algae.

**Ditch Mapping Unit**



Non-tidal, smaller waterways and graded or maintained permanent or temporary ditches that may or may not contain water at the time of the imagery.

**Open Water Mapping Unit**



Unchanneled tidal or non-tidal areas of standing water, including Suisun Bay.

**Developed Mapping Unit**




Includes parking lots, structures, water treatment ponds, and urban areas.

***Quercus agrifolia* Woodland Alliance**



*Quercus agrifolia* (coast live oak) is dominant or co-dominant in the tree canopy. Typically bordering freshwater creeks at upper reaches of marsh only, making this an uncommon vegetation type in Suisun Marsh.

***Eucalyptus (globulus, camaldulensis)* Woodland Semi-Natural Alliance**



*Eucalyptus camaldulensis*, *Eucalyptus globulus*, or other gum trees are dominant in the tree canopy. Planted as trees, groves, and windbreaks; naturalized on uplands and stream courses.

Understories are usually depauperate, though sometimes other non-natives, such as *Hedera helix*, clamber extensively in stands. Seeds of *Eucalyptus* germinate when tree crowns and built-up debris are removed by fire or in other ways.

***Salix laevigata* / *Salix lasiolepis* Association**



*Salix laevigata* (red willow) is dominant or co-dominant in the tree or shrub canopy with *S. lasiolepis* (Arroyo willow). Generally found at the edges of the marsh along freshwater creeks.



***Salix exigua* Association**



*Salix exigua* (narrow-leaf willow) is dominant or co-dominant in the shrub canopy with >20% absolute cover.

***Rosa californica* Association**



*Rosa californica* (California rose) is dominant and conspicuous, often forming narrow briar patches along levees and roads, occasionally in lower-lying portions of marsh.

***Rosa californica* – *Baccharis pilularis* Association**



*Rosa californica* (California rose) and *Baccharis pilularis* (coyote brush) co-occur in stand; either species may be dominant, but each has over 5% absolute cover. Often occurs along levees and roads.

***Baccharis salicifolia* Shrubland Alliance**




One stand strongly dominated by *Baccharis salicifolia* (mule-fat) occurs in the marsh; it was planted at the Blacklock restoration site along the southeastern levee.

***Arundo donax* Semi-Natural Association**



Clonal dense stands of *Arundo donax* (giant reed), generally small and locally distributed near settlements and roads in the marsh. The signature on the true-color imagery is slightly grayer than *Phragmites australis*, with which it might be confused. *Arundo* is also found in linear stands along roads or ditches more often than *Phragmites*, and the stand edges appear more ragged.

***Tamarix* spp. Shrubland Semi-Natural Alliance**



Stands are dominated by tamarisk, a large, non-native shrub with diffuse habit.

### Mediterranean Californian Naturalized Annual and Perennial Grassland Group




Upland grasslands generally are not associated with saturated soil or tidal influence throughout the growing season. Shrubs are generally less than 10% cover or, if more, sub-shrubs are overtopped by the dominant grass or herb species. Dominant species include the grasses *Hordeum murinum*, *Bromus* spp., *Festuca perennis*, *Elytrigia pontica*, and *Avena* spp. Also included in this group are stands dominated by weeds such as *Conium maculatum* (poison hemlock), *Foeniculum vulgare* (wild fennel), *Raphanus sativus* (wild radish), and *Brassica nigra* (black mustard).

### California Vernal Pool and Grassland Matrix Group



Vernal pools are temporary wetlands that fill with water during the rainy season and dry out in the spring.

### *Cortaderia (jubata, selloana)* Semi-Natural Herbaceous Alliance



Stands dominated by the large, non-native, mound-like *Cortaderia selloana* (pampas grass). Stands are generally small but conspicuous and occur in moist areas in the ecotone between wetlands and uplands. Some stands occur on the outboard sides of levees; however, all stands appear to be near buildings.




***Phalaris aquatica* Semi-Natural Association**



Stands are dominated by the tall bunch grass *Phalaris aquatica* (canary grass). These are usually small stands along levees but may occur in larger upland stands adjacent to the marsh.

***Rubus armeniacus* Shrubland Semi-Natural Alliance**



Vegetation is dominated by the introduced *Rubus armeniacus* (Himalayan berry), often in narrow briar patches along levees and roads in the Marsh. When co-dominant with *Rosa californica* or *Baccharis pilularis*, assign to those alliances.

***Baccharis pilularis* Shrubland Alliance**



*Baccharis pilularis* (coyote brush) is dominant to co-dominant with other shrubs such as *Atriplex lentiformis* (big saltbrush). *Rosa californica* (California rose) is typically absent or <<5% absolute cover.

***Baccharis pilularis* / Annual Grass – Herb Association**



*Baccharis pilularis* (coyote brush) dominates, with an understory that is typically dominated by annual grasses (*Hordeum* spp., *Festuca perennis*, *Bromus* spp.) and/or annual herbs.

***Carpobrotus edulis* or Other Ice Plants Herbaceous Semi-Natural Alliance**



Vegetation is dominated (>50% relative cover) by perennial non-native *Carpobrotus edulis* (iceplant), generally on levees and areas adjacent to buildings. Most stands are uniform and dense and have a distinct signature, particularly in the CIR imagery, where they are bright pink.

***Typha* (*angustifolia*, *latifolia*, *domingensis*) Herbaceous Alliance**



Stands are dominated by cattail species including *Typha angustifolia*, *T. latifolia*, and *T. domingensis* (Cattail (narrowleaf, broadleaf, and southern)). The distinguishing features of these three species are often blurred in the marsh and there is frequent evidence of hybridization. Different *Typha* species are often found in the same stand and are considered ecologically equivalent. Throughout most of the marsh, narrow-leaved forms (*T. angustifolia*/*domingensis*) predominate. Often, stands appear to be mostly dead-standing cattails from the previous season, but CIR imagery reveals that they are regenerating.



***Schoenoplectus (acutus, californicus)* Mapping Unit**



Vegetation dominated by *Schoenoplectus californicus* (California bulrush) and/or the ecologically and morphologically similar *Schoenoplectus acutus* (hardstem bulrush). Occasionally *Typha* spp. may occur in equal or higher cover than the *Schoenoplectus* spp., but *Schoenoplectus californicus* or *Schoenoplectus acutus* has at least 10% relative cover.

***Schoenoplectus (acutus, californicus)* / *Rosa californica* Association**



*Rosa californica* (California rose) is present with *Schoenoplectus californicus* (California bulrush), and/or *Schoenoplectus acutus* (hardstem bulrush). Usually found along levees bordering sloughs and channels, including the intertidal zone.

***Schoenoplectus (acutus, californicus)* – Wetland Herbs Mapping Unit**




Stands dominated by *Schoenoplectus californicus* (California bulrush) and/or *S. acutus* (hardstem bulrush) with an understory of >12% cover that is a varying mixture of mostly native perennial herbs, such as *Euthamia occidentalis*, *Aster lentus*, *A. subulatus*, *Artemisia douglasiana*, *Baccharis glutinosa*, *Achillea millefolium*, and *Stachys ajugoides*. May also include *Lepidium latifolium*.

***Schoenoplectus (acutus, californicus) – Typha (angustifolia, latifolia, domingensis) Mapping Unit***



Stands dominated in the overstory by *Schoenoplectus californicus* (California bulrush) and/or *Schoenoplectus acutus* (hardstem bulrush) with a lower to somewhat higher cover of *Typha angustifolia*, *T. latifolia*, and/or *T. domingensis*. May have up to 50% relative cover of wetland herbs (*Polygonum*, *Epilobium*, *Euthamia*, etc.).

***Phragmites australis* Herbaceous Alliance**



Most of the *Phragmites australis* (common reed) in the marsh is the very densely growing, invasive, non-native form. When *Phragmites* dominates, even when it is mixed with other plants such as *Schoenoplectus*, it is mapped as this type. The signature is very distinct, especially in the CIR imagery, in which it appears hot pink. There is no minimum mapping area for this type.

***Alisma triviale* Mapping Unit**



Vegetation is dominated by *Alisma triviale* (water plantain). Mapping is limited to a pond that receives freshwater drainage along Highway 680.



***Grindelia (stricta)* Provisional Herbaceous Alliance**




Stands dominated by the diffuse perennial herb *Grindelia* spp. (*Grindelia xpaludosa* may be more common in the marsh than previously thought and *Grindelia stricta* less so). May contain a variety of subordinate species, some weedy and some native. Typically found on the edges of wetlands on slightly elevated or drier ground than adjacent vegetation, such as natural or constructed levees, road margins, etc.

***Juncus arcticus* (var. *balticus*, *mexicanus*) Herbaceous Alliance**



Vegetation dominated or co-dominated (>30% relative cover) by the stoloniferous (clonal) rush *Juncus arcticus* var. *balticus* (Baltic rush) or *Juncus arcticus* var. *mexicanus* (Mexican rush), often associated with other taller or shorter herbaceous species. Usually in temporarily saturated wetlands not inundated for extensive periods. The signature on true color imagery is often very dark brown or black and its clonal habit is usually obvious.

***Leymus triticoides* Herbaceous Alliance**



Stands dominated (>50% relative cover) by the native *Leymus triticoides* (creeping wild rye). Stands are generally narrow bands of wetland-upland borders including natural ecotones between the *Distichlis spicata* (salt grass) Alliance and naturalized annual grass stands. Also occurs along levee tops and margins of marsh adjacent to vegetation of intermittent flooding zone. Non-natives, such as *Lepidium latifolium* and *Lotus corniculatus*, often invade these stands. Some very large stands, such as on Grizzly Island, may have been planted. The signature is bluer green in comparison to

***Lepidium latifolium* Herbaceous Semi-Natural Alliance**




Stands dominated by the invasive *Lepidium latifolium* (perennial pepperweed); may occur in temporarily flooded, intermittently flooded, and saturated wetlands, typically in at least slightly saline soils. Appears to be expanding in the marsh and is particularly threatening to native tidal marsh vegetation such as *Schoenoplectus americanus* (three-square), *Juncus arcticus* var. *balticus* (Baltic rush), and *Distichlis spicata* (saltgrass) Alliance stands (as at Rush Ranch). Often stands are at peak flower at the time of the imagery acquisition and so appear mounded and white. CIR imagery shows a very light pink signature.

***Persicaria lapathifolia* – *Xanthium strumarium* Herbaceous Alliance**



Vegetation of regularly disturbed, winter and vernal wet ponds and fields, usually on fine-grained clay-rich soils. *Xanthium strumarium* (cocklebur), *Persicaria lapathifolia* (common knotweed or willow weed), or other knotweed species are dominant or co-dominant in the herbaceous layer.

***Crypsis schoenoides* Mapping Unit**



Stands dominated by the low annual *Crypsis schoenoides* (swamp timothy). Found in winter and vernal flooded flats and pools. Vegetation is generally scattered, with intervening small to large openings of dry, cracked mud during summer.



### ***Salsola soda* Mapping Unit**



Stands dominated or co-dominated by *Salsola soda* (opposite-leaf Russian thistle). Large stands originated on the Hill Slough Unit.

### **Temperate and Boreal Salt Marsh Formation**



This type includes the former wetland herbs mapping unit and is applied when the vegetation is dominated by herbs in a wetland setting that does not have a recognizable signature or is too mixed in species composition to fit into a more specific vegetation type.

### ***Bolboschoenus maritimus* Herbaceous Alliance**



Vegetation of seasonally wet flats and pond bottoms dominated or co-dominated (>30% relative cover) by *Bolboschoenus maritimus* (alkali bulrush) in the taller herb/graminoid layer. Co-dominant species may include *Distichlis spicata*, *Salicornia pacifica*, *Sesuvium verrucosum*, *Spergularia salina*, *Typha latifolia*, and/or short herbs or grasses. Some stands also include the similar species *Bolboschoenus robustus* (sturdy bulrush) or hybrids between the two.

***Distichlis spicata* Herbaceous Alliance**



Stands are usually dominated (>50% relative cover) by *Distichlis spicata* (salt grass), or if not dominant, salt grass has higher cover than any other single species. *Distichlis spicata* has a grey-green or yellow-green signature in comparison to *Frankenia salina*, which is more blue-green and occurs in similar settings.

***Distichlis spicata* Association**



Stands strongly dominated by salt grass, *Distichlis spicata*, with no other species greater than 5% cover.

***Distichlis spicata* – Annual Grasses Association**



Stands are composed of a mixture of *Distichlis spicata* (salt grass) and non-native annual grasses. *Distichlis* may be dominant or share dominance (as low as 30% relative cover) with annual grass species, primarily *Polypogon monspeliensis* (rabbit's foot grass), *Festuca perennis* (perennial ryegrass), and/or *Hordeum* spp. (barley). Annuals generally cover at least 10%. Found at the higher marsh margins.



***Distichlis spicata* – *Juncus arcticus* var. *balticus* (*J. arcticus* var. *mexicanus*) Association**



Stands of *Distichlis spicata* (salt grass), with *Juncus arcticus* var. *balticus* or *Juncus arcticus* var. *mexicanus* as principal subordinate species (>5% relative cover). *Juncus* should be consistently present throughout the stand but does not need to be the most abundant subordinate species.

***Distichlis spicata* – *Salicornia pacifica* Association**



Stands are co-dominated by *Distichlis spicata* (salt grass) and *Salicornia pacifica* (pickleweed), both with 30% to 60% relative cover.

***Distichlis spicata* – *Schoenoplectus americanus* Provisional Association**




*Distichlis spicata* (salt grass) is the characteristic grass species with emergent *Schoenoplectus americanus* (three square bulrush) conspicuous, but less than 40% cover. Found in tidal and muted tidal settings with little disturbance.

***Distichlis spicata* – *Cotula coronopifolia* Association**



Stands of *Distichlis spicata* (salt grass) with the annual *Cotula coronopifolia* (brass buttons) as a subordinate species.

***Distichlis spicata* – *Juncus balticus* – *Triglochin* spp. – *Glaux maritima* Mapping Unit**



*Distichlis spicata* (salt grass) is the major ground cover, associated with a variety of native tidal marsh species including *Juncus balticus*, *Triglochin maritima*, *Glaux maritima*, *Jaumea carnosa*, and/or *Limonium californicum*. This type characterizes the most pristine native short tidal marsh habitats.

***Salicornia pacifica* (*Salicornia depressa*) Herbaceous Alliance**



Vegetation dominated (at least 10% cover over a sometimes higher cover of short annual or perennial grasses) by the native perennial salt marsh sub-shrubby or herbaceous *Salicornia pacifica* (pickleweed).



***Salicornia pacifica* Association**



Vegetation is dominated solely by *Salicornia pacifica* (pickleweed), with more than twice as much cover of *Salicornia* than of any other combination of species in the stand.

***Salicornia pacifica* – Annual Grasses Association**




Stand is dominated by *Salicornia pacifica* (pickleweed) with a sparse to dense mixture of annual grasses (*Polypogon*, *Hordeum*, *Festuca perennis*, *Bromus* spp.).

***Salicornia pacifica* – *Crypsis schoenoides* Association**




Vegetation dominated by *Salicornia pacifica* (pickleweed) mixed with a short intermittent layer of *Crypsis schoenoides* (swamp timothy).

***Salicornia pacifica* – *Sesuvium verrucosum* Association**



Vegetation dominated or co-dominated by *Salicornia pacifica* (pickleweed) with *Sesuvium verrucosum* (sea purslane) as a main subordinate species (at least 20% relative cover); may also include relatively high cover of *Cotula coronopifolia* (brass buttons).

***Salicornia pacifica* – *Cotula coronopifolia* Association**



Vegetation dominated by *Salicornia pacifica* with an ephemeral annual component of *Cotula coronopifolia* (brass buttons), which may cover enough ground to co-dominate in the early growing season.

***Spartina foliosa* Herbaceous Alliance**



*Spartina foliosa* was mapped along partially inundated open water along the bay.




***Atriplex prostrata* – *Cotula coronopifolia* Herbaceous Semi-Natural Alliance**



Stands dominated or characterized by *Atriplex prostrata* (fat hen) and/or *Cotula coronopifolia* (brass buttons). Both species are indicative of disturbed conditions in alkaline or saline wetlands. Both are early seral plants; they may be abundant to sparse from year to year depending on disturbance regime and salinity.

***Cotula coronopifolia* Semi-Natural Association**



Stands are strongly dominated by *Cotula coronopifolia* (brass buttons) with little or no significant cover from other species. Found on recently inundated, exposed mudflats. When still growing, the CIR imagery signature is a salmon color.

***Sesuvium verrucosum* Herbaceous Alliance**



*Sesuvium verrucosum* (sea purslane) > 50% relative cover in the herbaceous layer, dominant or co-dominant with *Chenopodium chenopodioides*, *Cotula coronopifolia*, *Distichlis spicata*, *Lolium perenne*, *Rumex crispus*, *Rumex pulcher*, *Salicornia pacifica* and/or *Spergularia marina*.

***Sesuvium verrucosum* Association**



*Sesuvium verrucosum* (sea purslane) is strongly dominant.

***Spergularia marina* Provisional Herbaceous Alliance**



Stands dominated by *Spergularia marina* (sand spurrey). Found in moist or seasonally flooded alkaline/saline areas, and newly scraped areas exposing mudflats of high salinity.

***Schoenoplectus americanus* Herbaceous Alliance**



*Schoenoplectus americanus* (three-square bulrush) dominates (>50% relative cover) stand. Co-dominants may include *Distichlis spicata*, *Juncus arcticus*, *Schoenoplectus acutus*, and *Schoenoplectus californicus*. *S. americanus* stands generally occupy portions of the marsh that are saturated, but not permanently flooded, often along the upper reaches of tidally influenced sloughs, creeks, and ditches.



***Schoenoplectus americanus* – *Lepidium latifolium* Association**



*Schoenoplectus americanus* (three-square bulrush) is dominant to co-dominant with *Lepidium latifolium* (perennial pepperweed), which may approach *S. americanus* in total cover. Tends to replace native associations of *S. americanus* along small tidal creeks and channels.

***Atriplex lentiformis* Shrubland Alliance**



Scrub dominated by the medium-to-large-sized shrub (up to 4 m in height) *Atriplex lentiformis* (big saltbrush). Generally, occurs in small, linear stands at borders of managed fields and intermittently flooded wetlands, usually associated with annual grasses and non-native herbs. In true color imagery, its signature is very gray in comparison to other shrubs of the same size.

***Frankenia salina* Herbaceous Alliance**



*Frankenia salina* (alkali heath) is dominant or co-dominant and may have equal or somewhat higher cover of *Distichlis spicata* (salt grass) or annual grasses. Generally found in seasonally moist or intermittently flooded clayey saline soils. *Frankenia* generally has a bluer green color relative to *Distichlis spicata*, which is more yellow-green but often has the same texture in appearance and occurs in similar settings.

***Frankenia salina* – *Distichlis spicata* Association**




*Frankenia salina* (alkali heath) and *Distichlis spicata* (saltgrass) co-dominate.

**Temperate Pacific Intertidal Flat Group**




This group-level mapping unit is used for floating aquatic vegetation that was not identifiable to species on the imagery and which had no field samples to verify its identification.

***Stuckenia pectinata* Association**



Floating masses strongly dominated by *Stuckenia pectinata* (narrow-leaved pondweed) occurring in diked brackish ponds, tidally influenced ditches or sloughs, and at the edges of the open bay.

### Row & Close Grain Crop Formation



Upland stands dominated by annual grasses and distinguished by heavily managed site history. Species are various but are planted, mowed, and/or cultivated regularly.

### Temperate Tree Developed Vegetation Group



Stands dominated by non-native or planted trees, except *Eucalyptus*, which are mapped as *Eucalyptus (globulus, camaldulensis)* Woodland Semi-Natural Alliance.



## Appendix C

### Map classes used to map Suisun Marsh

Vegetated map classes are ordered by their Group placement in the Manual of California Vegetation. 2021 map codes are given for each map class.

Group	Map Class (alliance/association)
Southwestern North American Introduced Riparian Scrub Group	
	Arundo donax Herbaceous Semi-Natural Association
	Tamarix spp. Shrubland Semi-Natural Alliance
Southwestern North American Salt Basin and High Marsh Group	
	Atriplex lentiformis Shrubland Alliance
Western North American Disturbed Alkaline Marsh and Meadow Group	
	Atriplex prostrata - Cotula coronopifolia Herbaceous Semi-Natural Alliance
	Cotula coronopifolia Semi-Natural Association
	Sesuvium verrucosum Herbaceous Alliance
	Sesuvium verrucosum Provisional Association
	Spergularia marina Provisional Herbaceous Alliance
California Coastal Evergreen Bluff and Dune Scrub Group	
	Baccharis pilularis / Annual Grass-Herb Association
	Baccharis pilularis Shrubland Alliance
Southwestern North American Riparian/Wash Scrub Group	
	Baccharis salicifolia Shrubland Alliance
Temperate Pacific Tidal Salt and Brackish Meadow Group	
	Bolboschoenus maritimus Herbaceous Alliance
	Distichlis spicata - Annual grasses Association
	Distichlis spicata - Cotula coronopifolia Association
	Distichlis spicata - Juncus arcticus var. balticus (J. arcticus var. mexicanus) Association
	Distichlis spicata - Juncus balticus - Triglochin spp. - Glaux maritima Mapping Unit
	Distichlis spicata - Salicornia pacifica Association
	Distichlis spicata - Schoenoplectus americanus Provisional Association
	Distichlis spicata Association
	Distichlis spicata Herbaceous Alliance
	Salicornia pacifica - Annual Grasses Association
	Salicornia pacifica - Cotula coronopifolia Association
	Salicornia pacifica - Crypsis schoenoides Association
	Salicornia pacifica - Sesuvium verrucosum Association
	Salicornia pacifica (Salicornia depressa) Herbaceous Alliance
	Salicornia pacifica Association
	Spartina foliosa Herbaceous Alliance
California-Vancouverian Semi-Natural Littoral Scrub and Herb Vegetation Group	
	Carpobrotus edulis or Other Ice Plants Herbaceous Semi-Natural Alliance
Mediterranean California Naturalized Annual and Perennial Grassland Group	
	Mediterranean California Naturalized Annual and Perennial Grassland Group
	Cortaderia (jubata, selloana) Semi-Natural Herbaceous Alliance



Introduced North American Mediterranean Woodland and Forest Macrogroup (no subdivision at group level)	
	Eucalyptus (globulus, camaldulensis) Woodland Semi-Natural Alliance
Southwestern North American Salt Basin and High Marsh Group	
	Frankenia salina - Distichlis spicata Association
	Frankenia salina Herbaceous Alliance
Californian Mixed Annual/Perennial Freshwater Vernal Pool/Swale Bottomland Group	
	Grindelia (stricta) Provisional Herbaceous Alliance
California Vernal Pool and Grassland Matrix Group	
	California Vernal Pool and Grassland Matrix Group
Californian Warm Temperate Marsh/Seep Group	
	Juncus arcticus (var. balticus, mexicanus) Herbaceous Alliance
	Leymus triticoides Alliance
Naturalized Warm-Temperate Riparian and Wetland Group	
	Crypsis schoenoides Mapping Unit
	Lepidium latifolium Herbaceous Semi-Natural Alliance
	Persicaria lapathifolia - Xanthium strumarium Herbaceous Alliance
	Salsola soda Mapping Unit
Vancouverian and Rocky Mountain Naturalized Perennial Grassland Group	
	Phalaris aquatica Herbaceous Semi-Natural Association
California Broadleaf Forest and Woodland	
	Quercus agrifolia Woodland Alliance
Row & Close Grain Crop Formation	
	Row and Close Grain Crop Formation
Naturalized Non-Native Deciduous Scrub Group	
	Rubus armeniacus Shrubland Semi-Natural Alliance
Southwestern North American Riparian/Wash Scrub Group	
	Salix exigua Association
	Rosa californica Association
	Rosa californica - Baccharis pilularis Association
Southwestern North American Riparian Evergreen and Deciduous Woodland Group	
	Salix laevigata / Salix lasiolepis Association
Southwestern North American Alkali Marsh/Seep Vegetation Group	
	Schoenoplectus americanus - Lepidium latifolium Association
	Schoenoplectus americanus Herbaceous Alliance
Temperate and Boreal Salt Marsh Formation	
	Temperate and Boreal Salt Marsh Formation
Temperate Pacific Intertidal Flat Group	
	Stuckenia pectinata Association
	Temperate Pacific Intertidal Flat Group
Temperate Tree Developed Vegetation Group	
	Temperate Tree Developed Vegetation Group
Arid West Freshwater Emergent Marsh Group	
	Typha (angustifolia, latifolia, domingensis) Herbaceous Alliance
	Schoenoplectus (acutus, californicus) - Rosa californica Association

	Schoenoplectus (acutus, californicus) - Wetland Herbs Mapping Unit
	Schoenoplectus (acutus, californicus) - Typha (angustifolia, latifolia, domingensis) Mapping Unit
	Schoenoplectus (acutus, californicus) Mapping Unit
	Phragmites australis Herbaceous Alliance
	Alisma triviale Mapping Unit
Non-Vegetation	
	Bare Ground Mapping Unit
	Developed Mapping Unit
	Ditch Mapping Unit
	Open Water Mapping Unit
	Railroad Track Mapping Unit
	Road / Trails Mapping Unit
	Slough Mapping Unit
	Tidal Mudflat Mapping Unit

## Appendix D

### 2018 Vegetation Mapping Attributes and Descriptions

**OBJECTID\_1:** Internal feature number

**MapClass:** The finest level of vegetation type mapped (alliance, association, group, macrogroup, or mapping unit); or land use for polygons that are not natural vegetation.

**MapCode:** Numeric code corresponding to the MapClass.

**PER\_HARDWOOD:** The total cover of hardwood tree species present in the stand. "0.2" represents hardwood presence at less than 1%.

**PER\_CONIFER:** The total cover of conifer tree species present in the stand. "0.2" represents conifer presence at less than 1%.

**PER\_TREE:** The sum of % hardwood cover and % conifer cover.

**PER\_SHRUB:** The total cover of all shrub species present.

**HERB\_CODE:** The cover of herbaceous species present in a stand, broken into the following classes: <2%, 2-9%, 10-39%, >40% Woody (used when PER\_TREE+PER\_SHRUB is >40%).

**PER\_TOTAL:** The total combined cover of % tree + % shrub + average of the herbaceous cover by class

**HT\_CODE:** Average height of the dominant vegetation within the polygon, in meters, broken into the following classes: <1 m, 1-5 m, 5-20 m, 20-50 m

**SIZE\_CATEGORY:** Used only for tree mapping types, the average diameter at breast height of the trees present within the stand, broken into the following classes: Saplings (1-6"), Pole (6-11"), Small (11-24"), Medium-Large (>24").

**ISOLATED\_TREE:** A "Y" indicates a total tree cover of >0% and <5%.

**RESTORATION:** A "Y" indicates obvious signs of restoration present in the polygon.

**CLEARING\_DISTURBANCE:** The percent of the polygon that has been affected by human disturbance (including disking, plowing, and mowing), broken into the following classes:  
No Disturbance = <5%, Minimal Disturbance = 5-25%, Moderate Disturbance = 25-50%, High Disturbance = >50%

**INVASIVE\_PLANT:** The percent of the polygon that is covered by invasive species, broken into the following classes:  
No Invasive = <5%, Low Invasive = 5-25%, Moderate Invasive = 25-50%, High Invasive = >50%

**Comments\_Final:** Comments from the photo interpreter about an individual polygon

**Habitat:**

Tidal (1): Tidal wetlands (including muted tidal) are those areas naturally affected regularly by tidal fluctuation. The area may or may not be vegetated with vascular or non-vascular plants and may or

may not have any evidence of human modification such as ditches, excavations, interrupted levees, berms, etc.

Leveed (2): leveed wetlands are those areas that are completely enclosed and are restricted from natural tidal influence.

Slough (3): Generally applied to the wide, naturally occurring (at least not linear) waterways affected by tides.

**Unique\_ID:** Unique identifier for each polygon.

**MCVName:** Names will be the same as the ones used in the Manual of California Vegetation and California Natural Community List. These standardized names correspond as much as possible to the National Vegetation Classification System. Classes for human land use or otherwise unvegetated land were drawn from the California Wildlife Habitat Relationship.

**MCVLevel:** The standardized level of the vegetation description in the Manual of California Vegetation corresponding with levels of the National Vegetation Classification System.

**CalVegType:** A crosswalk to the CalVeg vegetation system. Note that there may be a one-to-many relationship between CalVeg and NVCS.

**CalVegCode:** The two-letter code for the crosswalked CalVeg vegetation system. Note that there may be a one-to-many relationship between CalVeg and NVCS.

**CWHRType:** A crosswalk to the California Wildlife Habitat Relationships (CWHR) system. Note that there is usually a one-to-many relationship between CWHR and NVCS.

**CWHRCode:** The three-letter code for the crosswalked CWHR habitat type. Note that there is usually a one-to-many relationship between CWHR and NVCS.

**GlobalRank:** The global rarity rank of the plant community (only for polygons mapped to the alliance level).

- G1: Fewer than 6 viable occurrences and/or 2000 acres worldwide;
- G2: 6-20 viable occurrences and/or 2000-10,000 acres worldwide;
- G3: 21-100 viable occurrences and/or 10,000-50,000 acres worldwide;
- G4: Greater than 100 viable occurrences and/or greater than 50,000 acres worldwide;
- G5: Community demonstrably secure due to secure worldwide abundance.

**StateRank:** The state rarity rank of the plant community (only for polygons mapped to the alliance level).

- S1: Fewer than 6 viable occurrences and/or 2000 acres statewide;
- S2: 6-20 viable occurrences and/or 2000-10,000 acres statewide;
- S3: 21-100 viable occurrences and/or 10,000-50,000 acres statewide;
- S4: Greater than 100 viable occurrences and/or greater than 50,000 acres statewide;
- S5: Community demonstrably secure due to secure statewide abundance.

**Sensitive:** Rarity of the vegetation type. Alliances and associations with state ranks of S1-S3 and global ranks of G1-G3 are considered sensitive.

**Y** – Sensitive

**N** – Not Sensitive

**CaCode:** California Natural Community Codes - unique code assigned to alliances and associations.

**MCVAlliance:** The Manual of California Vegetation (MCV)-compliant alliance (if applicable) which contains this vegetation type.

**MCVGroup:** The MCV-compliant group (if applicable) which contains this vegetation type.

**MCVMacrogroup:** The MCV-compliant macrogroup (if applicable) which contains this vegetation type.

**Community\_Link:** The link to the alliance in the Manual of California Vegetation Online.

**Ortho\_NAIP\_Year:** All imagery for this mapping effort was flown in 2018.

**Shape\_Length:** Length of the polygon, in meters.

**Shape\_Area:** Area of the polygon, in meters squared.

**Acres:** The area of a polygon in acres

**Hectares:** The area of a polygon in hectares

## Appendix E

### Salt Marsh Harvest Mouse habitat by vegetation mapping type

Note: some mapping types at group level and higher could not be rated as potential habitat.

Mapping Code	Mapping type	Potential Habitat
<b>1</b>	Bare Ground Mapping Unit	No
<b>2</b>	Agriculture	No
<b>4</b>	Road / Trails Mapping Unit	No
<b>6</b>	Slough Mapping Unit	No
<b>7</b>	Tidal Mudflat Mapping Unit	No
<b>8</b>	Railroad Track Mapping Unit	No
<b>9</b>	Ditch Mapping Unit	No
<b>11</b>	Open Water Mapping Unit	No
<b>15</b>	Developed Mapping Unit	No
<b>100000</b>	<b>Warm Temperate Forest Formation</b>	No
<b>110000</b>	<b>Madrean Forest and Woodland Division</b>	No
<b>111000</b>	<b>California Forest and Woodland Macrogroup</b>	No
1111000	Californian Broadleaf Forest and Woodland Group	No
1111100	<i>Quercus agrifolia</i> Alliance	No
1111101	<i>Quercus agrifolia</i> Association	No
1111102	<i>Quercus agrifolia</i> / <i>Salix lasiolepis</i> Association	No
1111200	<i>Quercus lobata</i> Alliance	No
<b>200000</b>	<b>Cool Temperate Forest Formation</b>	No
<b>210000</b>	<b>North American Introduced Evergreen Broadleaf and Conifer Forest Division</b>	No
<b>211000</b>	<b>Introduced North American Mediterranean Woodland and Forest Macrogroup</b>	No
2110100	<i>Eucalyptus (globulus, camaldulensis)</i> Semi-Natural Alliance	No
2110101	<i>Eucalyptus globulus</i> Semi-Natural Association	No
2110200	<i>Ailanthus altissima</i> Semi-Natural Alliance	No
<b>212000</b>	<b>Southwestern North American Riparian, Flooded and Swamp Forest Macrogroup</b>	No
2121000	Southwestern North American Riparian Evergreen and Deciduous Woodland Group	No
2121100	<i>Salix laevigata</i> Alliance	No
2121101	<i>Salix laevigata</i> / <i>Salix lasiolepis</i> Association	No
2122000	Southwestern North American Riparian/Wash Scrub Group	Not rated
2122100	<i>Salix exigua</i> Alliance	No
2122101	<i>Salix exigua</i> Association	No
2122200	<i>Rosa californica</i> Alliance	Yes
2122210	<i>Rosa californica</i> Association	Yes
2122212	<i>Rosa californica</i> – <i>Baccharis pilularis</i> Association	Yes
2122300	<i>Baccharis salicifolia</i> Shrubland Alliance	Yes

Mapping Code	Mapping type	Potential Habitat
2123000	Southwestern North American Introduced Riparian Scrub Group	No
2123100	<i>Arundo donax</i> Semi-Natural Alliance	No
2123101	<i>Arundo donax</i> Semi-Natural Association	No
2123200	<i>Tamarix</i> spp. Shrubland Semi-Natural Alliance	No
<b>3000000</b>	<b>Western North America Flooded and Swamp Forest Division</b>	No
<b>3100000</b>	<b>Western Cordilleran montane–boreal riparian scrub Macrogroup</b>	No
3111000	Vancouverian Riparian Deciduous Forest Group	No
3111100	<i>Fraxinus latifolia</i> Alliance	No
3111101	<i>Fraxinus latifolia</i> Planted Stands Mapping Unit	No
<b>4000000</b>	<b>Mediterranean Grassland and Forb Meadow Formation</b>	Not rated
<b>4100000</b>	<b>California Grassland and Meadow Division</b>	Not rated
<b>4110000</b>	<b>California Annual and Perennial Grassland Macrogroup</b>	Not rated
4111000	Mediterranean California Naturalized Annual and Perennial Grassland Group	Yes
4111100	<i>Brassica nigra</i> and Other Mustards Semi-Natural Alliance	No
4111101	<i>Brassica nigra</i> Semi-Natural Association	No
4111102	<i>Raphanus sativus</i> Semi-Natural Association	No
4111200	<i>Bromus (diandrus, hordeaceus) – Brachypodium distachyon</i> Semi-Natural Alliance	Yes
4111300	<i>Centaurea (solstitialis, melitensis)</i> Semi-Natural Alliance	No
4111400	<i>Conium maculatum – Foeniculum vulgare</i> Semi-Natural Alliance	No
4111401	<i>Conium maculatum</i> Semi-Natural Association	No
4111402	<i>Foeniculum vulgare</i> Semi-Natural Association	No
4111500	<i>Cortaderia (jubata, selloana)</i> Semi-Natural Herbaceous Alliance	No
4111600	<i>Festuca perennis</i> Semi-Natural Alliance	Yes
4111601	<i>Festuca perennis – Lepidium latifolium</i> Semi-Natural Association	Yes
4111602	<i>Festuca perennis – Lotus corniculatus</i> Semi-Natural Association	Yes
4111603	<i>Festuca perennis – Rumex</i> spp. Mapping Unit	Yes
4111604	<i>Hordeum marinum – Festuca perennis</i> Mapping Unit	Yes
4111700	<i>Elytrigia pontica</i> Mapping Unit	Yes
4111800	<i>Agrostis avenacea</i> Mapping Unit	Yes
4111900	<i>Vulpia</i> spp. – <i>Euthamia occidentalis</i> Mapping Unit	Yes
<b>5000000</b>	<b>Temperate Grassland, Meadow, and Shrubland Formation</b>	Not rated
<b>5100000</b>	<b>Vancouverian and Rocky Mountain Grassland and Shrubland Division</b>	Not rated
<b>5110000</b>	<b>Western North American Temperate Grassland and Meadow Macrogroup</b>	Not rated
5111000	Vancouverian and Rocky Mountain Naturalized Perennial Grassland Group	Not rated
5111100	<i>Phalaris aquatica</i> Herbaceous Semi-Natural Alliance	Yes

Mapping Code	Mapping type	Potential Habitat
5111101	<i>Phalaris aquatica</i> Herbaceous Semi-Natural Association	Yes
<b>5120000</b>	<b>Vancouverian Lowland Grassland and Shrubland Macrogroup</b>	Not rated
5121000	Naturalized Non-Native Deciduous Scrub Group	Not rated
5121100	<i>Rubus armeniacus</i> Shrubland Semi-Natural Alliance	Yes
<b>6000000</b>	<b>Temperate and Boreal Scrub and Herb Coastal Vegetation Formation</b>	Not rated
<b>6100000</b>	<b>Pacific Coast Scrub and Herb Littoral Vegetation Division</b>	Not rated
<b>6110000</b>	<b>Vancouverian Coastal Dune and Bluff Macrogroup</b>	Not rated
6111000	California Coastal Evergreen Bluff and Dune Scrub Group	Not rated
6111100	<i>Baccharis pilularis</i> Shrubland Alliance	Yes
6111101	<i>Baccharis pilularis</i> / Annual Grass-Herb Association	Yes
6112000	California–Vancouverian Semi-Natural Littoral Scrub and Herb Vegetation Group	No
6112100	<i>Carpobrotus edulis</i> or Other Ice Plants Herbaceous Semi-Natural Alliance	No
<b>7000000</b>	<b>Temperate and Boreal Freshwater Marsh Formation</b>	Not rated
<b>7100000</b>	<b>Western North American Freshwater Marsh Division</b>	Not rated
<b>7110000</b>	<b>Western North American Freshwater Marsh Macrogroup</b>	Not rated
7111000	Arid West Freshwater Emergent Marsh Group	Not rated
7111100	<i>Typha (angustifolia, domingensis, latifolia)</i> Herbaceous Alliance	No
7111101	<i>Typha (angustifolia, latifolia, domingensis)</i> Association	No
7111102	<i>Typha (angustifolia, latifolia, domingensis)</i> (dead stalks) Mapping Unit	No
7111103	<i>Typha (angustifolia, latifolia, domingensis)</i> – <i>Echinochloa crus-galli</i> Association	No
7111104	<i>Typha (angustifolia, latifolia, domingensis)</i> – <i>Distichlis spicata</i> Association	No
7111105	<i>Typha (angustifolia, latifolia, domingensis)</i> – <i>Phragmites australis</i> Association	Yes
7111106	<i>Typha (angustifolia, latifolia, domingensis)</i> – <i>Schoenoplectus americanus</i> Association	Yes
7111200	<i>Schoenoplectus (acutus, californicus)</i> Alliance	Yes
7111201	<i>Schoenoplectus californicus</i> – <i>Schoenoplectus acutus</i> Association	Yes
7111202	<i>Schoenoplectus (acutus, californicus)</i> Mapping Unit	Yes
7111203	<i>Schoenoplectus (acutus, californicus)</i> / <i>Rosa californica</i> Provisional Association	Yes
7111300	<i>Schoenoplectus (acutus, californicus)</i> – <i>Typha (angustifolia, latifolia, domingensis)</i> Mapping Unit	Yes
7111400	<i>Schoenoplectus (acutus, californicus)</i> – Wetland Herbs Mapping Unit	Yes
7111401	<i>Calystegia sepium</i> – <i>Euthamia occidentalis</i> Mapping Unit	No
7111500	<i>Phragmites australis</i> Herbaceous Alliance	Yes
7111501	<i>Phragmites australis</i> Association	Yes



Mapping Code	Mapping type	Potential Habitat
7111502	<i>Phragmites australis</i> – <i>Schoenoplectus</i> spp. Association	Yes
7111503	<i>Phragmites australis</i> – <i>Xanthium strumarium</i> Mapping Unit	No
7111600	<i>Alisma triviale</i> Mapping Unit	No
7112000	Vancouverian Coastal/Tidal Marsh and Meadow Group	Not rated
7112100	<i>Potentilla anserina</i> Alliance	Yes
<b>7120000</b>	<b>Western North America Vernal Pool Macrogroup</b>	Not rated
7121000	Californian Mixed Annual/Perennial Freshwater Vernal Pool/Swale Bottomland Group	Not rated
7121000	California Vernal Pool and Grassland Matrix Group	Yes
7121100	<i>Grindelia (camporum, stricta)</i> Provisional Herbaceous Alliance	Yes
<b>7130000</b>	<b>Western North America Wet Meadow and Low Shrub Carr Macrogroup</b>	Not rated
7131000	Californian Warm Temperate Marsh/Seep Group	Not rated
7131100	<i>Juncus arcticus</i> (var. <i>balticus</i> , <i>mexicanus</i> ) Herbaceous Alliance	Yes
7131101	<i>Juncus arcticus</i> var. <i>balticus</i> Association	Yes
7131102	<i>Juncus arcticus</i> var. <i>balticus</i> – <i>Conium maculatum</i> Association	Yes
7131103	<i>Juncus arcticus</i> var. <i>balticus</i> – <i>Lepidium latifolium</i> Association	Yes
7131104	<i>Juncus arcticus</i> var. <i>balticus</i> – <i>Potentilla anserina</i> Association	Yes
7131200	<i>Leymus triticoides</i> Alliance	Yes
7132000	Naturalized Warm-Temperate Riparian And Wetland Group	Yes
7132100	<i>Lepidium latifolium</i> Herbaceous Semi-Natural Alliance	Yes
7132200	<i>Persicaria lapathifolia</i> – <i>Xanthium strumarium</i> Provisional Herbaceous Alliance	Yes
7132201	<i>Polygonum</i> spp. – <i>Xanthium strumarium</i> – <i>Echinochloa crus-galli</i> Mapping Unit	Yes
7132300	<i>Cynodon dactylon</i> – <i>Crypsis</i> spp. – <i>Paspalum</i> spp. Moist Semi-Natural Alliance	Yes
7132301	<i>Crypsis schoenoides</i> Mapping Unit	Yes
7132302	<i>Cynodon dactylon</i> Mapping Unit	No
7132400	<i>Polypogon monspeliensis</i> Mapping Unit	Yes
7132500	<i>Rumex</i> spp. Mapping Unit	Yes
7132600	<i>Salsola soda</i> Mapping Unit	No
<b>8000000</b>	<b>Temperate and Boreal Salt Marsh Formation</b>	Not rated
<b>8100000</b>	<b>Temperate and Boreal Pacific Coastal Salt Marsh Division</b>	Not rated
<b>8110000</b>	<b>North American Pacific Coastal Salt Marsh Macrogroup</b>	Not rated
8111000	Temperate Pacific Tidal Salt and Brackish Meadow Group	Not rated
8111100	<i>Bolboschoenus maritimus</i> Herbaceous Alliance	Yes
8111101	<i>Bolboschoenus maritimus</i> Association	Yes
8111102	<i>Bolboschoenus maritimus</i> – <i>Salicornia pacifica</i> Association	Yes

Mapping Code	Mapping type	Potential Habitat
8111103	<i>Bolboschoenus maritimus</i> – <i>Sesuvium verrucosum</i> Association	Yes
8111200	<i>Distichlis spicata</i> Herbaceous Alliance	Yes
8111201	<i>Distichlis spicata</i> Association	Yes
8111202	<i>Distichlis spicata</i> – Annual Grasses Association	Yes
8111203	<i>Distichlis spicata</i> – <i>Juncus arcticus</i> var. <i>balticus</i> ( <i>J. arcticus</i> var. <i>mexicanus</i> ) Association	Yes
8111204	<i>Distichlis spicata</i> – <i>Lotus corniculatus</i> Mapping Unit	No
8111205	<i>Distichlis spicata</i> – <i>Salicornia pacifica</i> Association	Yes
8111206	<i>Distichlis spicata</i> – <i>Schoenoplectus americanus</i> Provisional Association	Yes
8111207	<i>Distichlis spicata</i> – <i>Cotula coronopifolia</i> Association	Yes
8111208	<i>Distichlis spicata</i> – <i>Bolboschoenus maritimus</i> Mapping Unit	Yes
8111209	<i>Distichlis spicata</i> – <i>Juncus balticus</i> – <i>Triglochin</i> spp. – <i>Glaux maritima</i> Mapping Unit	Yes
8111210	<i>Lepidium latifolium</i> – <i>Distichlis spicata</i> Semi-Natural Association	Yes
8111300	<i>Salicornia pacifica</i> ( <i>Salicornia depressa</i> ) Herbaceous Alliance	Yes
8111301	<i>Salicornia pacifica</i> Tidal Association	Yes
8111302	<i>Salicornia pacifica</i> – Annual Grasses Association	Yes
8111303	<i>Salicornia pacifica</i> – <i>Atriplex prostrata</i> Association	Yes
8111304	<i>Salicornia pacifica</i> – <i>Crypsis schoenoides</i> Association	Yes
8111305	<i>Salicornia pacifica</i> – <i>Sesuvium verrucosum</i> Association	Yes
8111306	<i>Salicornia pacifica</i> – <i>Echinochloa crus-galli</i> – <i>Polygonum</i> – <i>Xanthium strumarium</i> Association	Yes
8111307	<i>Salicornia pacifica</i> – <i>Cotula coronopifolia</i> Association	Yes
8111400	<i>Spartina foliosa</i> Herbaceous Alliance	Yes
8112000	Western North American Disturbed Alkaline Marsh and Meadow Group	Not rated
8112100	<i>Atriplex prostrata</i> – <i>Cotula coronopifolia</i> Herbaceous Semi-Natural Alliance	Yes
8112101	<i>Cotula coronopifolia</i> Semi-Natural Association	Yes
8112102	<i>Atriplex prostrata</i> Semi-Natural Association	Yes
8112103	<i>Atriplex prostrata</i> – <i>Distichlis spicata</i> Semi-Natural Association	Yes
8112104	<i>Atriplex prostrata</i> – <i>Bolboschoenus maritimus</i> Semi-Natural Association	Yes
8112105	<i>Atriplex prostrata</i> – <i>Sesuvium verrucosum</i> Semi-Natural Association	Yes
8112106	<i>Atriplex prostrata</i> – Annual Grasses Semi-Natural Association	Yes
8112200	<i>Sesuvium verrucosum</i> Herbaceous Alliance	Yes
8112201	<i>Sesuvium verrucosum</i> Association	Yes
8112202	<i>Sesuvium verrucosum</i> – <i>Distichlis spicata</i> Association	Yes
8112203	<i>Sesuvium verrucosum</i> – <i>Festuca perennis</i> Association	Yes
8112204	<i>Sesuvium verrucosum</i> – <i>Cotula coronopifolia</i> Association	Yes
8112300	<i>Spergularia marina</i> Provisional Herbaceous Alliance	No
8112301	<i>Spergularia marina</i> – <i>Cotula coronopifolia</i> Mapping Unit	No
8200000	Western North American Interior Alkali–Saline Wetland Division	Not rated
8210000	Warm Semi-Desert/Mediterranean Alkali–Saline Wetland Macrogroup	Not rated

Mapping Code	Mapping type	Potential Habitat
8211000	Southwestern North American Alkali Marsh/Seep Vegetation Group	Not rated
8211100	<i>Schoenoplectus americanus</i> Herbaceous Alliance	Yes
8211101	<i>Schoenoplectus americanus</i> Association	Yes
8211102	<i>Schoenoplectus americanus</i> – <i>Potentilla anserina</i> Association	Yes
8211103	<i>Schoenoplectus americanus</i> – <i>Schoenoplectus californicus</i> – <i>Schoenoplectus acutus</i> Association	Yes
8211104	<i>Schoenoplectus americanus</i> – <i>Lepidium latifolium</i> Association	Yes
8212000	Southwestern North American Salt Basin and High Marsh Group	Not rated
8212100	<i>Atriplex lentiformis</i> Shrubland Alliance	Yes
8212200	<i>Frankenia salina</i> Herbaceous Alliance	Yes
8212201	<i>Frankenia salina</i> – Annual grasses Mapping Unit	Yes
8212202	<i>Frankenia salina</i> – <i>Distichlis spicata</i> Association	Yes
<b>9000000</b>	<b>Hydromorphic Vegetation (Aquatic Vegetation) Class</b>	No
<b>9100000</b>	<b>Marine and Estuarine Saltwater Aquatic Vegetation Formation</b>	No
<b>9110000</b>	<b>Temperate Pacific Saltwater Aquatic Vegetation Division</b>	No
<b>9111000</b>	<b>Temperate Pacific Intertidal Shore Macrogroup</b>	No
9111100	Temperate Pacific Intertidal Flat Group	No
9111200	<i>Stuckenia (pectinata)</i> – <i>Potamogeton</i> spp. Alliance	No
9111201	<i>Stuckenia pectinata</i> Association	No
9111300	<i>Ruppia (cirrhosa, maritima)</i> Herbaceous Alliance	No
<b>9200000</b>	<b>North American Freshwater Aquatic Vegetation Division</b>	No
<b>9210000</b>	<b>Western North American Freshwater Aquatic Vegetation Macrogroup</b>	No
9211000	Naturalized Temperate Pacific Freshwater Vegetation Group	No
9211100	<i>Ludwigia (hexapetala, peploides)</i> Provisional Herbaceous Semi-Natural Alliance	No
<b>9310000</b>	<b>Herbaceous Agricultural Vegetation Subclass</b>	No
<b>9311000</b>	<b>Row &amp; Close Grain Crop Formation</b>	No
<b>9312000</b>	<b>Fallow Field and Weed Vegetation Formation</b>	No
<b>9312100</b>	<b>Cropland Fallow Field Division</b>	No
<b>9312110</b>	<b>Fallow Field Macrogroup</b>	No
<b>9410000</b>	<b>Herbaceous &amp; Woody Developed Vegetation Subclass</b>	No
<b>9411000</b>	<b>Other Developed Vegetation Formation</b>	No
<b>9411100</b>	<b>Other Developed Vegetation Division</b>	No
9411111	Temperate Tree Developed Vegetation Group	No

## Appendix F

Acreage and acreage change of Salt Marsh Harvest Mouse habitat in Suisun Marsh within the tidal, leveed, and whole region, within the four management regions and marsh-wide in 1999, 2015, 2018, and 2021.

Potential SMHM Habitat	Tidal									
	Acres				Change in Acres			Percent Change		
	1999	2015	2018	2021	1999-2021	2015-2018	2018-2021	1999-2021	2015-2018	2018-2021
Management Region 0	N/A	N/A	79.08	169.65	N/A	N/A	90.57	N/A	N/A	114.53%
Management Region 1	1529.3	2329.6	2683.9	2,682.17	1,152.87	354.3	-1.73	75.39%	15.21%	-0.06%
Management Region 2	1360.8	2088.6	2137.1	2,160.90	800.10	48.5	23.8	58.80%	2.32%	1.11%
Management Region 3	448.9	597.3	652	757.2	308.30	54.7	105.2	68.68%	9.16%	16.13%
Management Region 4	2316.5	2643.9	3258.3	3,236.60	920.10	614.4	-21.7	39.72%	23.24%	-0.67%
Marsh-wide	5655.5	7659.4	8810.3	9,006.52	3,351.02	1150.9	196.22	59.25%	15.03%	2.23%

Potential SMHM Habitat	Leveed									
	Acres				Change in Acres			Percent Change		
	1999	2015	2018	2021	1999-2021	2015-2018	2018-2021	1999-2021	2015-2018	2018-2021
Management Region 0	N/A	N/A	2.99	3.7	N/A	N/A	0.71	N/A	N/A	23.75%
Management Region 1	11758.5	8545	10033.7	9,388.90	-2,369.60	1488.7	-644.8	-20.15%	17.42%	-6.43%
Management Region 2	5558.4	5321.9	5975.4	5,474.20	-84.20	653.5	-501.2	-1.51%	12.28%	-8.39%
Management Region 3	3114.1	2375.6	2654.9	2,456.40	-657.70	279.3	-198.5	-21.12%	11.76%	-7.48%
Management Region 4	21018.8	19780.7	21628.7	20,307.30	-711.50	1848	-1321.4	-3.39%	9.34%	-6.11%
Marsh-wide	41449.8	36023.1	40295.7	37,655.50	-3,794.30	4272.6	-2640.2	-9.15%	11.86%	-6.55%

Potential SMHM Habitat	Total									
	Acres				Change in Acres			Percent Change		
	1999	2015	2018	2021	1999-2021	2015-2018	2018-2021	1999-2021	2015-2018	2018-2021
Management Region 0	N/A	N/A	82.07	173.4	N/A	N/A	91.33	N/A	N/A	111.28%
Management Region 1	13287.8	10874.6	12707.6	12,071.10	-1,216.70	1833	-636.5	-9.16%	16.86%	-5.01%
Management Region 2	6919.3	7410.5	8112.5	7,635.10	715.80	702	-477.4	10.34%	9.47%	-5.88%
Management Region 3	3562.9	2972.9	3306.8	3,213.60	-349.30	333.9	-93.2	-9.80%	11.23%	-2.82%
Management Region 4	23335.4	22424.5	24887	23,543.90	208.50	2462.5	-1343.1	0.89%	10.98%	-5.40%
Marsh-wide	47105.3	43682.5	49106	46,662	-443.30	5423.5	-2444	-0.94%	12.42%	-4.98%

## Appendix G

Acreage and acreage change of the non-native species of concern in Suisun Marsh within the tidal, leveed, and whole region, within the four management regions and marsh-wide in 1999, 2015, 2018, and 2021.

### Management Region 0

Species	Tidal									
	Acres				Change in Acres			Percent Change		
	1999	2015	2018	2021	1999-2018	2015-2018	2018-2021	1999-2018	2015-2018	2018-2021
Management Region 0										
<i>Arundo donax</i> Association	N/A	N/A	0.3	0.31	N/A	N/A	0.01	N/A	N/A	3.33%
<i>Carpobrotus edulis</i> Association	N/A	N/A	0	0.04	N/A	N/A	0.04	N/A	N/A	N/A
<i>Cortaderia selloana</i> Association	N/A	N/A	0.02	0.01	N/A	N/A	-0.01	N/A	N/A	-62.87%
<i>Eucalyptus</i> spp.	N/A	N/A	0.21	0.42	N/A	N/A	0.22	N/A	N/A	105.36%
<i>Lepidium latifolium</i> *	N/A	N/A	0.08	0.01	N/A	N/A	-0.07	N/A	N/A	-87.50%
<i>Phragmites australis</i>	N/A	N/A	9.01	13.61	N/A	N/A	4.60	N/A	N/A	51.00%
<i>Salsola soda</i> Provisional Alliance	N/A	N/A	0	0	N/A	N/A	0	N/A	N/A	N/A

Species	Leveed									
	Acres				Change in Acres			Percent Change		
	1999	2015	2018	2021	1999-2018	2015-2018	2018-2021	1999-2018	2015-2018	2018-2021
Management Region 0										
<i>Arundo donax</i> Association	N/A	N/A	0.05	0.06	N/A	N/A	0.01	N/A	N/A	20.55%
<i>Carpobrotus edulis</i> Association	N/A	N/A	0	0	N/A	N/A	0	N/A	N/A	N/A
<i>Cortaderia selloana</i> Association	N/A	N/A	0	0	N/A	N/A	0	N/A	N/A	N/A
<i>Eucalyptus</i> spp.	N/A	N/A	3.21	3.53	N/A	N/A	0.32	N/A	N/A	9.89%
<i>Lepidium latifolium</i> *	N/A	N/A	0.01	0.05	N/A	N/A	0.04	N/A	N/A	363.68%
<i>Phragmites australis</i>	N/A	N/A	0.34	0.53	N/A	N/A	0.19	N/A	N/A	55.30%
<i>Salsola soda</i> Provisional Alliance	0	29.9	0	0	N/A	N/A	0	N/A	N/A	N/A

Species	Total									
	Acres				Change in Acres			Percent Change		
	1999	2015	2018	2021	1999-2018	2015-2018	2018-2021	1999-2018	2015-2018	2018-2021
Management Region 0										
<i>Arundo donax</i> Association	N/A	N/A	0.35	0.37	N/A	N/A	0.02	N/A	N/A	5.93%
<i>Carpobrotus edulis</i> Association	N/A	N/A	0	0.04	N/A	N/A	0.04	N/A	N/A	N/A
<i>Cortaderia selloana</i> Association	N/A	N/A	0.02	0.01	N/A	N/A	-0.01	N/A	N/A	-50.00%
<i>Eucalyptus</i> spp.	N/A	N/A	3.42	3.95	N/A	N/A	0.53	N/A	N/A	15.62%
<i>Lepidium latifolium</i> *	N/A	N/A	0.09	0.06	N/A	N/A	-0.03	N/A	N/A	-30.15%
<i>Phragmites australis</i>	N/A	N/A	9.35	14.14	N/A	N/A	4.79	N/A	N/A	51.16%
<i>Salsola soda</i> Provisional Alliance	0	29.9	0	0	N/A	N/A	0	N/A	N/A	N/A



**Management Region 1**

Species	Tidal									
	Acres				Change in Acres			Percent Change		
	1999	2015	2018	2021	1999-2018	2015-2018	2018-2021	1999-2018	2015-2018	2018-2021
<b>Management Region 1</b>										
<i>Arundo donax</i> Association	0	2.2	1	2.34	2.34	-1.2	1.34	N/A	-54.55%	57.26%
<i>Carpobrotus edulis</i> Association	0	1.9	0	0	0.00	-1.9	0	N/A	-100.00%	N/A
<i>Cortaderia selloana</i> Association	0	0.3	1	0.48	0.48	0.7	-0.52	N/A	233.33%	-106.94%
<i>Eucalyptus</i> spp.	0	0.9	0.3	0.79	0.79	-0.6	0.49	N/A	-66.67%	61.84%
<i>Lepidium latifolium</i> *	31.6	7	14.35	7.32	-24.28	7.35	-7.03	-76.82%	105.00%	-95.93%
<i>Phragmites australis</i>	102.8	454.6	587.2	589.55	486.75	132.6	2.35	473.50%	29.17%	0.40%
<i>Salsola soda</i> Provisional Alliance	0	0	0	0	0.00	0	0	N/A	N/A	N/A

Species	Leveed									
	Acres				Change in Acres			Percent Change		
	1999	2015	2018	2021	1999-2018	2015-2018	2018-2021	1999-2018	2015-2018	2018-2021
<b>Management Region 1</b>										
<i>Arundo donax</i> Association	0.7	0	1.8	2.09	1.39	1.8	0.29	198.57%	N/A	16.11%
<i>Carpobrotus edulis</i> Association	0	0	2.5	2.16	2.16	2.5	-0.34	N/A	N/A	-13.46%
<i>Cortaderia selloana</i> Association	7.6	3.6	5.2	3.84	-3.76	1.6	-1.36	-49.44%	44.44%	-26.11%
<i>Eucalyptus</i> spp.	52.6	91	102.6	101.31	48.71	11.6	-1.29	92.61%	12.75%	-1.26%
<i>Lepidium latifolium</i> *	159	20.2	34.6	35.81	-123.19	14.4	1.21	-77.48%	71.29%	3.51%
<i>Phragmites australis</i>	82.4	162.4	307.8	389.53	307.13	145.4	81.73	372.73%	89.53%	26.55%
<i>Salsola soda</i> Provisional Alliance	0	29.9	13.9	26.32	26.32	-16	12.42	N/A	-53.51%	89.32%

Species	Total									
	Acres				Change in Acres			Percent Change		
	1999	2015	2018	2021	1999-2018	2015-2018	2018-2021	1999-2018	2015-2018	2018-2021
<b>Management Region 1</b>										
<i>Arundo donax</i> Association	0.7	2.2	2.8	4.43	3.73	0.6	1.63	532.86%	27.27%	58.21%
<i>Carpobrotus edulis</i> Association	0	1.9	2.5	2.16	2.16	0.6	-0.34	N/A	31.58%	-13.46%
<i>Cortaderia selloana</i> Association	7.6	3.9	6.2	4.33	-3.27	2.3	-1.87	-43.08%	58.97%	-30.23%
<i>Eucalyptus</i> spp.	52.6	91.8	103	102.10	49.50	11.2	-0.90	94.10%	12.20%	-0.88%
<i>Lepidium latifolium</i> *	190.6	27.2	49	43.14	-147.46	21.8	-5.86	-77.37%	80.15%	-11.96%
<i>Phragmites australis</i>	185.2	617.1	895	979.09	793.89	277.9	84.09	428.67%	45.03%	9.40%
<i>Salsola soda</i> Provisional Alliance	0	29.9	13.9	26.32	26.32	-16	12.42	N/A	-53.51%	89.32%

**Management Region 2**

Species	Tidal									
	Acres				Change in Acres			Percent Change		
	1999	2015	2018	2021	1999-2018	2015-2018	2018-2021	1999-2018	2015-2018	2018-2021
<b>Management Region 2</b>										
<i>Arundo donax</i> Association	0.2	0	0	0	-0.2	0	0	-100.00%	N/A	N/A
<i>Carpobrotus edulis</i> Association	0	0	0	0	0	0	0	N/A	N/A	N/A
<i>Cortaderia selloana</i> Association	0.8	0	0.02	0.03	-0.77	0.02	0.01	-96.25%	N/A	50.00%
<i>Eucalyptus</i> spp.	3.2	2.2	2	6.16	2.96	-0.2	4.16	92.50%	-9.09%	208.00%
<i>Lepidium latifolium</i> *	52.9	9.5	14.9	11.99	-40.91	5.4	-2.91	-77.33%	-9.09%	-19.53%
<i>Phragmites australis</i>	17.2	125.8	183.2	235.03	217.83	57.4	51.83	1266.45%	44.12%	28.29%
<i>Salsola soda</i> Provisional Alliance	0	0	0	0	0	0	0	N/A	N/A	N/A

Species	Leveed									
	Acres				Change in Acres			Percent Change		
	1999	2015	2018	2021	1999-2018	2015-2018	2018-2021	1999-2018	2015-2018	2018-2021
<b>Management Region 2</b>										
<i>Arundo donax</i> Association	0	0	0.5	0.38	0.38	0.5	-0.12	N/A	N/A	-23.78%
<i>Carpobrotus edulis</i> Association	0	0.4	1.4	1.24	1.24	1	-0.16	N/A	250%	-11.53%
<i>Cortaderia selloana</i> Association	0	0	0	0	0.00	0	0.00	N/A	N/A	N/A
<i>Eucalyptus</i> spp.	29.7	27	26.8	21.14	-8.56	-0.2	-5.66	-28.82%	-0.74%	-21.11%
<i>Lepidium latifolium</i> *	120.3	4.6	30.1	31.30	-89.00	25.5	1.20	-73.99%	554.35%	3.97%
<i>Phragmites australis</i>	53.6	226.1	311.9	316.44	262.84	85.8	4.54	490.38%	37.95%	1.46%
<i>Salsola soda</i> Provisional Alliance	0	8.2	0	0	0.00	-8.2	0.00	N/A	-100%	N/A

Species	Total									
	Acres				Change in Acres			Percent Change		
	1999	2015	2018	2021	1999-2018	2015-2018	2018-2021	1999-2018	2015-2018	2018-2021
<b>Management Region 2</b>										
<i>Arundo donax</i> Association	0.2	0	0.5	0.38	0.18	0.5	-0.12	91%	N/A	-23.78%
<i>Carpobrotus edulis</i> Association	0	0.4	1.4	1.24	1.24	1	-0.16	N/A	250%	-11.53%
<i>Cortaderia selloana</i> Association	0.8	0	0.02	0.03	-0.77	0.02	0.01	-96%	N/A	50.00%
<i>Eucalyptus</i> spp.	32.8	29.2	28.9	27.30	-5.50	-0.3	-1.60	-17%	-0.34%	-5.53%
<i>Lepidium latifolium</i> *	173.3	14.1	45	43.29	-130.01	30.9	-1.71	-75%	205.40%	-3.81%
<i>Phragmites australis</i>	70.8	351.9	495.1	551.47	480.67	143.2	56.37	679%	39.90%	11.39%
<i>Salsola soda</i> Provisional Alliance	0	8.2	0	0.00	0.00	-8.2	0.00	N/A	-100%	N/A

**Management Region 3**

Species	Tidal									
	Acres				Change in Acres			Percent Change		
	1999	2015	2018	2021	1999-2018	2015-2018	2018-2021	1999-2018	2015-2018	2018-2021
<b>Management Region 3</b>										
<i>Arundo donax</i> Association	0.6	0	0	0	-0.6	0	0	-100%	N/A	N/A
<i>Carpobrotus edulis</i> Association	0	0	0	0	0	0	0	N/A	N/A	N/A
<i>Cortaderia selloana</i> Association	0	0	0	0	0	0	0	N/A	N/A	N/A
<i>Eucalyptus</i> spp.	0	0.2	0.36	0.41	0.41	0.16	0.05	N/A	80.00%	13.89%
<i>Lepidium latifolium</i> *	0	0	0	0.03	0.03	0	0.03	N/A	N/A	N/A
<i>Phragmites australis</i>	1.2	31.6	100.6	131.35	130.15	69	30.75	10845.83%	218.35%	30.57%
<i>Salsola soda</i> Provisional Alliance	0	0	0	0	0	0	0	N/A	N/A	N/A

Species	Leveed									
	Acres				Change in Acres			Percent Change		
	1999	2015	2018	2021	1999-2018	2015-2018	2018-2021	1999-2018	2015-2018	2018-2021
<b>Management Region 3</b>										
<i>Arundo donax</i> Association	0	0	0	0	0	0	0	N/A	N/A	N/A
<i>Carpobrotus edulis</i> Association	0	0	0	0	0	0	0	N/A	N/A	N/A
<i>Cortaderia selloana</i> Association	0	0	0	0	0	0	0	N/A	N/A	N/A
<i>Eucalyptus</i> spp.	1.6	7.1	7.2	7.11	5.51	0.1	-0.09	344.38%	1.41%	-1.25%
<i>Lepidium latifolium</i> *	0	0	5.8	5.9	5.9	5.8	0.1	N/A	N/A	1.72%
<i>Phragmites australis</i>	25.1	137	135.5	94.05	68.95	-1.5	-41.45	274.70%	50.33%	-30.59%
<i>Salsola soda</i> Provisional Alliance	0	0	0	0	0	0	0	N/A	N/A	N/A

Species	Total									
	Acres				Change in Acres			Percent Change		
	1999	2015	2018	2021	1999-2018	2015-2018	2018-2021	1999-2018	2015-2018	2018-2021
<b>Management Region 3</b>										
<i>Arundo donax</i> Association	0.6	0	0	0	-0.6	0	0	-100%	N/A	N/A
<i>Carpobrotus edulis</i> Association	0	0	0	0	0	0	0	N/A	N/A	N/A
<i>Cortaderia selloana</i> Association	0	0	0	0	0	0	0	N/A	N/A	N/A
<i>Eucalyptus</i> spp.	1.6	7.3	7.5	7.52	5.92	0.2	0.02	370.00%	2.74%	0.27%
<i>Lepidium latifolium</i> *	0	0	5.8	5.93	5.93	5.8	0.13	N/A	N/A	2.24%
<i>Phragmites australis</i>	26.3	168.5	236.1	225.4	199.1	67.6	-10.7	757.03%	40.12%	-4.53%
<i>Salsola soda</i> Provisional Alliance	0	0	0	0	0	0	0	N/A	N/A	N/A

**Management Region 4**

Species	Tidal									
	Acres				Change in Acres			Percent Change		
	1999	2015	2018	2021	1999-2018	2015-2018	2018-2021	1999-2018	2015-2018	2018-2021
<b>Management Region 4</b>										
<i>Arundo donax</i> Association	1.1	0.1	2.2	1.66	1.1	2.1	-0.54	100.00%	2200%	-24.55%
<i>Carpobrotus edulis</i> Association	1.6	0.1	0	0.23	-1.6	-0.1	0.23	-100.00%	-100.00%	N/A
<i>Cortaderia selloana</i> Association	1.2	2.2	2.8	0.09	1.6	0.6	-2.71	133.33%	27.27%	-96.79%
<i>Eucalyptus</i> spp.	9.5	4.5	4.2	6.11	-5.3	-0.3	1.91	-55.79%	-6.67%	45.48%
<i>Lepidium latifolium</i> *	18.7	0	3.3	3.05	-15.4	3.3	-0.25	-82.35%	N/A	-7.58%
<i>Phragmites australis</i>	240.4	570.8	984.8	941.23	744.4	414	-43.57	309.65%	72.53%	-4.42%
<i>Salsola soda</i> Provisional Alliance	0	0	0	0	0	0	0	N/A	N/A	N/A

Species	Leveed									
	Acres				Change in Acres			Percent Change		
	1999	2015	2018	2021	1999-2018	2015-2018	2018-2021	1999-2018	2015-2018	2018-2021
<b>Management Region 4</b>										
<i>Arundo donax</i> Association	2.1	3.7	4.3	2.49	0.39	0.6	-1.81	18.57%	16.22%	-42.09%
<i>Carpobrotus edulis</i> Association	5.5	30.7	36.2	36.56	31.06	5.5	0.36	564.73%	17.92%	0.99%
<i>Cortaderia selloana</i> Association	0.2	0.2	0.7	0	-0.2	0.5	-0.7	-100.00%	250.00%	-100.00%
<i>Eucalyptus</i> spp.	111.6	156	173	173.17	61.57	17	0.17	55.17%	10.90%	0.10%
<i>Lepidium latifolium</i> *	263.8	66.7	102.5	149.6	-114.2	35.8	47.1	-43.29%	53.67%	45.95%
<i>Phragmites australis</i>	170.4	1238.6	1675.4	1,729.66	1559.26	436.8	54.26	915.06%	35.27%	3.24%
<i>Salsola soda</i> Provisional Alliance	0	28.9	25.2	42.05	42.05	-3.7	16.85	N/A	-12.80%	66.87%

Species	Total									
	Acres				Change in Acres			Percent Change		
	1999	2015	2018	2021	1999-2018	2015-2018	2018-2021	1999-2018	2015-2018	2018-2021
<b>Management Region 4</b>										
<i>Arundo donax</i> Association	3.3	3.8	6.4	4.15	0.85	2.6	-2.25	25.76%	68.42%	-35.16%
<i>Carpobrotus edulis</i> Association	7	30.8	36.2	36.79	29.79	5.4	0.59	425.57%	17.53%	1.63%
<i>Cortaderia selloana</i> Association	1.4	2.4	3.5	0.09	-1.31	1.1	-3.41	-93.57%	45.83%	-97.43%
<i>Eucalyptus</i> spp.	121.1	160.5	177.2	179.28	58.18	16.7	2.08	48.04%	10.40%	1.17%
<i>Lepidium latifolium</i> *	282.5	66.7	105.8	152.65	-129.85	39.1	46.85	-45.96%	58.62%	44.28%
<i>Phragmites australis</i>	410.8	1809.4	2660.2	2670.89	2260.09	850.8	10.69	550.17%	47.02%	0.40%
<i>Salsola soda</i> Provisional Alliance	0	28.9	25.2	42.05	42.05	-3.7	16.85	N/A	-12.80%	66.87%

**Marsh-wide**

Species	Tidal									
	Acres				Change in Acres			Percent Change		
	1999	2015	2018	2021	1999-2018	2015-2018	2018-2021	1999-2018	2015-2018	2018-2021
<b>Marsh-wide</b>										
<i>Arundo donax</i> Association	1.9	2.3	3.5	4.32	2.42	1.2	0.82	127%	52.17%	23.43%
<i>Carpobrotus edulis</i> Association	1.6	2	0	0.27	-1.33	-2	0.27	-83%	-100.00%	N/A
<i>Cortaderia selloana</i> Association	2	2.5	3.85	0.61	-1.39	1.35	-3.24	-70%	54.00%	-84.16%
<i>Eucalyptus</i> spp.	12.7	7.7	7.1	13.89	1.19	-0.6	6.79	9%	-7.79%	95.63%
<i>Lepidium latifolium</i> *	103.3	16.5	32.6	22.4	-80.9	16.1	-10.2	-78%	97.58%	-31.29%
<i>Phragmites australis</i>	361.6	1182.8	1864.7	1910.77	1549.17	681.9	46.07	428%	57.65%	2.47%
<i>Salsola soda</i> Provisional Alliance	0	0	0	0	0	0	0	N/A	N/A	N/A

Species	Leveed									
	Acres				Change in Acres			Percent Change		
	1999	2015	2018	2021	1999-2018	2015-2018	2018-2021	1999-2018	2015-2018	2018-2021
<b>Marsh-wide</b>										
<i>Arundo donax</i> Association	2.8	3.7	6.6	5.03	2.23	2.9	-1.57	79.64%	78.38%	-23.79%
<i>Carpobrotus edulis</i> Association	5.5	31	40.2	39.96	34.46	9.2	-0.24	626.55%	29.68%	-0.60%
<i>Cortaderia selloana</i> Association	7.8	3.8	5.85	3.84	-3.96	2.05	-2.01	-50.77%	53.95%	-34.36%
<i>Eucalyptus</i> spp.	195.4	281.1	312.9	306.27	110.87	31.8	-6.63	56.74%	11.31%	-2.12%
<i>Lepidium latifolium</i> *	543.1	91.5	173.1	222.66	-320.44	81.6	49.56	-59.00%	89.18%	28.63%
<i>Phragmites australis</i>	331.6	1764.2	2431.1	2530.22	2198.62	666.9	99.12	663.03%	37.80%	4.08%
<i>Salsola soda</i> Provisional Alliance	0	67	39.1	68.36	68.36	-27.9	29.26	N/A	-41.64%	74.83%

Species	Total									
	Acres				Change in Acres			Percent Change		
	1999	2015	2018	2021	1999-2018	2015-2018	2018-2021	1999-2018	2015-2018	2018-2021
<b>Marsh-wide</b>										
<i>Arundo donax</i> Association	4.7	6.1	10.1	9.35	4.65	4	-0.75	114.89%	65.57%	-7.43%
<i>Carpobrotus edulis</i> Association	7	33.1	40.2	40.23	33.23	7.1	0.03	474.29%	21.45%	0.07%
<i>Cortaderia selloana</i> Association	9.8	6.3	9.7	4.45	-5.35	3.4	-5.25	-1.02%	53.97%	-54.12%
<i>Eucalyptus</i> spp.	208.1	288.8	320	320.16	112.06	31.2	0.16	53.77%	10.80%	0.05%
<i>Lepidium latifolium</i> *	646.4	108	205.7	245.06	-401.34	97.7	39.36	-68.18%	90.46%	19.13%
<i>Phragmites australis</i>	693.1	2947	4295.8	4440.99	3747.89	1348.8	145.19	519.80%	45.77%	3.38%
<i>Salsola soda</i> Provisional Alliance	0	67	39.1	68.36	68.36	-27.9	29.26	N/A	-41.64%	74.83%



## RECON FIELD FORM - SUISUN MARSH (May 27, 2015)

# 2021 Vegetation Map Update for Suisun Marsh

1-2	2-5	>5										
Field Alliance/Association name:												
Comments:												
% Cover: Tree _____ Shrub _____ Herb _____ Total Veg _____ Veg Ht (m): <0.5 0.5-1 1-2 2-5 5-10 >10 Disturb: N/A Low Med High												
Strata	Species	% cover		Strata	Species	% cover		Strata	Species	% cover		

CALIFORNIA DEPARTMENT OF FISH AND WILDLIFE PROTOCOL FOR  
RECONNAISSANCE FIELD FORM

**Definitions of fields in the form**

**I. LOCATIONAL/ENVIRONMENTAL DESCRIPTION**

**Recorder:** The full name of the recorder should be provided for the first field form for the day. On successive forms, initials can be recorded.

**Other Surveyors:** The full name of each person assisting should be provided for the first field form for the day. On successive forms, initials of each person assisting can be recorded.

**Date:** Date of the sampling.

**Return?:** Check this box if team members should return to this spot at a later date to take a recon or RA/relevé. This can be used if the phenology is not conducive to identification of the major species, or if there is not enough time to take the survey.

**Waypoint ID:** The waypoint number assigned by a Global Positioning System (GPS) unit when marking and storing a waypoint for the sample location.

**UID:** The ID number of a reference point or polygon that this reconnaissance describes.

**Location Name:** The name of the property, park, or location within large holdings (like USFS or BLM properties).

**GPS name:** The name/number assigned to the GPS unit.

**Projected? Yes / No / Base / Digitized:** Circle the appropriate option:

**Yes** - The point is a projected, or offset point. The surveyor used a bearing and distance to project the point to match what they are describing with the survey.

**No** - The surveyors are in the vegetation they are describing and the point is where the observer was standing for photographs. This location can also be used as a base location for an offset survey.

**Base** - Base point only. This is where a surveyor was standing when taking an offset survey to describe vegetation not at that point. No plant data or vegetation descriptions are associated with this location. However, cardinal photos taken at this point will be stored in a directory of this name.

**Digitized** – An offset point was created on the GPS unit without taking bearing and distance readings. This option should only be used when the imagery on the GPS unit is unique and unmistakable.

**Bearing (°):** The compass bearing from the Base point to the Projected point.

**Distance (m):** The distance in meters from the Base point to the Projected point, determined by use of a range finder.

**Inclination (°):** The vertical offset from the Base point to the Projected point.

**Base Waypoint ID:** For a projected or digitized point, this is the location where the surveyor was standing when the information was collected. Cardinal photographs will be taken at this point and will


be stored on the computer under this ID. Photographs of the stand vegetation will be taken from this point and will be stored on the computer under the Projected point's ID.

**Base / Projected UTM's:** If the point is projected or digitized, circle whether the coordinates of the base point or the offset point have been recorded. These will generally be for the offset point.

**PDOP:** The accuracy of the GPS location. Record the error reading from the GPS unit.

**GPS coordinates:** Record easting (**UTME**) and northing (**UTMN**) from a GPS unit.

**Stand Size:** Estimate the size of the entire stand in which the sample is taken and circle the appropriate range. As a measure, one acre is similar in size to a football field.

**Camera/Photos:** Write the name camera, JPG numbers, and direction of the photos. Take four photos in the main cardinal directions (N, E, S, W) clockwise from the north, from the GPS location. This symbol can be used to indicate the cardinal photos: . If additional photos are taken in other directions, please note the JPG numbers and a description of each photo.

## II. HABITAT AND VEGETATION DESCRIPTION

**Field Alliance/Association name:** Name of alliance or association following the most recent Suisun Marsh classification, using scientific nomenclature, *e.g.*, *Quercus agrifolia*. An alliance is based on the dominant or diagnostic species of the stand and usually reflects the uppermost and/or dominant height stratum. A dominant species covers the greatest area. A diagnostic species is consistently found in some vegetation types but not others.

Please note: The field-assessed alliance name may not exist in the present classification, in which case you can provide a new alliance name in this field.

**Comments:** Briefly describe the stand age/seral stage, disturbance history, nature and extent of land use, and other site environmental and vegetation factors that will aid in the mapping effort.

### % Cover:

**Tree:** The total cover of all the trees taking into consideration the porosity, or the holes, in the vegetation. This is an estimate of the absolute tree cover, disregarding the overlap<sup>1</sup> of individual trees.

**Shrub:** The total cover of all the shrubs taking into consideration the porosity, or the holes, in the vegetation. This is an estimate of the absolute shrub cover, disregarding the overlap of individual shrubs.

**Herb:** The total cover of all the herbs taking into consideration the porosity, or the holes, in the vegetation. This is an estimate of the absolute herbaceous cover, disregarding the overlap<sup>1</sup> of individual herbs.

**Total Veg:** The total cover of all vascular vegetation taking into consideration the porosity, or the holes, in the vegetation. This is an estimate of the absolute vegetation cover, disregarding the overlap<sup>1</sup> of the various tree, shrub, and/or herbaceous layers and species.

**Veg Ht (m):** Modal height for all vegetation. Estimate the mean height and circle the appropriate height range.

**Disturb.:** Estimate the amount of disturbance in the stand from human activity, such as roads, trails, disking, tilling, clearing, etc.

**N/A** = not applicable for this polygon type

**Low** = 0-33% of polygon affected by disturbance

**Med** = 34-66% of polygon affected by disturbance

**High** = >67% of polygon affected by disturbance

### Species List and Coverage

List the species that are dominant or that are characteristically consistent throughout the stand. This list is used if there is some uncertainty in the field-assessed alliance name, so the most common species should be listed. In the interests of time and efficiency, this species list should not be exhaustive.

#### Strata:

**T = Tree.** A woody perennial plant that has a single trunk.

**A = Sapling.** 1" - <6" dbh and young in age, OR small trees that are <1" dbh, are clearly of appreciable age and are kept short by repeated browsing, burning, or other disturbance. Includes trees that are re-sprouting from roots or stumps following fire, logging, or other disturbance. These re-sprouts may exhibit a shrubby form, with multiple small trunks, but are species that are generally considered trees. If a majority of the trunks are >6" dbh, then the re-sprouts would be recorded under the "Tree" stratum.

**E = Seedling.** A tree species clearly of a very young age that is < 1" dbh or has not reached breast height. This applies only to trees propagating from seed; re-sprouts are not recorded here even if they meet the size requirements.

**S = Shrub.** A perennial, woody plant, that is multi-branched and doesn't die back to the ground every year.

**H = Herb.** An annual or perennial that dies down to ground level every year.

**N = Non-vascular.** Includes moss, lichen, liverworts, hornworts, cryptogamic crust, and algae.

When one or more tree species are regenerating, the Tree, Seedling, and/or Sapling strata may be noted on the same line, e.g.:

Strata	Species	%Cover	C
T/A/E	Quercus douglasii	40/<1/<1	C

**Species:** Use Jepson Manual nomenclature. When uncertain of an identification (which you intend to confirm later) use parentheses to indicate what part of the determination needs to be confirmed. For example, you could write out *Brassica (nigra)* if you are sure it is a *Brassica* but you need further clarification on the specific epithet.

**% cover:** provide the % absolute aerial cover for each species listed. All species percent covers may total over 100% because of overlap.

**Collections:** If a species collection is made, it should be indicated in the blank column next to "% cover" with a "C" (for collected). If the species is later keyed out, cross out the species name or description and write the keyed species name in pen on the datasheet. Do not erase what was written in the field, because this information can be used if specimens get mixed up later. If the specimen is then thrown out, add a "T" to the "C" in that column (CT = thrown out after confirmation) or cross out the "C". If the specimen is kept but is still not confidently identified, add a "U" to the "C" (CU = collected and unconfirmed). In this case, the unconfirmed species epithet should be put in parentheses [e.g. *Hordeum (murinum)*]. If the specimen is kept and is confidently identified, add a "C" to the existing "C" (CC =



collected and confirmed). If the specimen is later deposited in an herbarium, add a "D" to the existing "C" (CD = collected and deposited) and note the receiving herbarium.