

Table SPECIES. Species number taken by hook-and-line, set line, stick gear and miscellaneous sampled in the Big Sur Commerical Nearshore Skiff Fishery, 1997-2001.

<u>Species</u>			
<u>Rank</u>	<u>Common Name</u>	<u>No. Taken</u>	<u>% Composition</u>
1	Cabazon	2213	32.98%
2	Black-and-yellow rf.	1538	22.92%
3	Gopher rf.	1085	16.17%
4	Grass rf.	744	11.09%
5	Blue rf.	254	3.78%
6	Vermilion rf.	183	2.73%
7	Olive rf.	124	1.85%
8	Kelp greenling	94	1.40%
9	Lingcod	90	1.34%
10	Yellowtail rf.	69	1.03%
11	Kelp rf.	42	0.63%
12	Copper rf.	41	0.61%
13	Bocaccio	37	0.55%
14	Widow rf.	30	0.45%
15	Greenspotted rf.	28	0.42%
16	Starry rf.	20	0.30%
17	Canary rf.	17	0.25%
17	Chilipepper	17	0.25%
18	Speckled rf.	14	0.21%
19	Black rf.	13	0.19%
19	Unidentified rf.	13	0.19%
20	Rock greenling	12	0.18%
21	Rosy rf.	11	0.16%
22	California sheephead	8	0.12%
23	Greenstriped rf.	5	0.07%
24	Brown rf.	3	0.04%
25	China rf.	2	0.03%
26	Ocean whitefish	1	0.01%
26	Pacific bonito	1	0.01%
26	Pacific mackerel	1	0.01%
26	Yelloweye rf.	1	0.01%